

THE GEOMETRY OF HIDDEN TREE MARKOV MODELS FOR BINARY DATA

PIOTR ZWIERNIK AND JIM Q. SMITH

ABSTRACT. In this paper we investigate the geometry of a discrete Bayesian network whose graph is a tree all of whose variables are binary and the only observed variables are those labeling its leaves. We obtain a full geometric description of these models which is given by polynomial equations and inequalities. Our analysis is based on combinatorial results generalizing the notion of cumulants so that they apply to the models under analysis. The geometric structure we obtain links to the notion of a tree metric considered in phylogenetic analysis and to some interesting determinantal formulas involving the hyperdeterminant of $2 \times 2 \times 2$ tables.

1. INTRODUCTION

A Bayesian network whose graph is a tree all of whose inner nodes represent variables which are not directly observed lie in an important class of models, containing phylogenetic tree models and hidden Markov models. Inference for this model class tends to be challenging and often needs to employ fragile numerical algorithms. In [27] we established a useful new coordinate system for such models when all of the variables are binary. This analysis enabled us not only to address various identifiability issues but also helped us to derive exact formulas for the maximum likelihood estimators given that the sample proportions were consistent with edge probabilities assigned to this model class.


However, the application of this new coordinate system reaches far beyond understanding the identifiability and it can be used to analyze the global structure of these tree models. For example [7] gave an intriguing correspondence between, on the one hand, a correlation system on tree models and on the other distances induced by trees where the length between two nodes in a tree is given as a sum of the length of edges in the path joining them. Our new coordinate system for the tree models enables us to explore this relationship between probabilistic tree models and tree metrics in detail. It was already implicit that constraints on possible distances between any two leaves in the tree imply some inequality constraints on the possible covariances between the binary variables represented by the leaves. These inequalities follow from the four-point condition ([21], Definition 7.1.5) together with some other simple non-negativity constraints (c.f. Equation (27)). However in this paper we also show that these inequality constraints cannot be sufficient and there are some additional constraints involving higher order tree cumulants. We provide the full set of the defining constraints in Theorem 4.6. This is given by a list

Key words and phrases. conditional independence, graphical models on trees, general Markov models, hidden data, binary data, central moments, cumulants, Möbius function, semi-algebraic statistical models, inequality constraints, phylogenetic invariants, tree metrics.

of polynomial equations and inequalities which describe the set of all probability distributions in the model.

Our approach here is founded in a geometric study of tree models through the method of phylogenetic invariants first introduced by Lake [15], and Cavender and Felsenstein [6]. These invariant algebraic relationships are expressed as a set of polynomial equations over the observed probability tables which must hold for a given phylogenetic model to be valid. We note that these algebraic techniques have also been embraced by computational algebraic geometers [1][11][26] enhancing the statistical and computational analyses of such models [5]. A similar problems can be solved for other model classes [8]. The main technical deficiency of using phylogenetic invariants in this way is that they do not give a *full* geometric description of the statistical model. The additional inequalities obtained as the main result of this paper complete this description. Where and how these inequality constraints can helpfully supplement an analysis based on phylogenetic invariants is illustrated by the simple example given below.

Example 1.1. Let T be the tripod tree below where we use the convention that observed nodes are depicted by shaded nodes



The inner node represents a binary hidden variable H and the leaves represent binary observable variables X_1, X_2, X_3 . The tree represents the conditional independence statements $X_1 \perp\!\!\!\perp X_2 \perp\!\!\!\perp X_3 | H$. The model has full dimension over the observed margin (X_1, X_2, X_3) and consequently there are no equations defining it. However it is not a saturated model since not all the marginal probability distributions over the observed vector (X_1, X_2, X_3) lie in the model. For example Lazarsfeld [16, Section 3.1] showed that the second moments of the observed distribution must satisfy

$$\text{Cov}(X_1, X_2)\text{Cov}(X_1, X_3)\text{Cov}(X_2, X_3) \geq 0.$$

This constraint, which clearly impacts the inferences we might want to make, is not acknowledged through the study of phylogenetic invariants. Therefore inference based solely on these invariants is incomplete and in particular naive estimates derived through these methods can be infeasible within the model class in a sense illustrated later in this paper.

This example motivated the closer investigation of the semi-algebraic features associated with the geometry of binary tree models with hidden inner nodes. The main problem with the geometric analysis of these models is that in general it is hard to obtain the inequality constraints defining a model even for very simple examples (see [9, Section 4.3][12, Section 7]). Despite this, some results can be found in the literature. Thus in the case of a binary naive Bayes model a somewhat complicated solution was given by Auvray et al. [2]. In the binary case there are also some partial results for general tree structures given by Pearl and Tarsi [19] and Steel and Faller [25]. The most important applications in biology involve variables that can take four values. Recently Matsen [17] gave a set of inequalities in this case for

group-based phylogenetic models (additional symmetries are assumed) using the Fourier transformation of the raw probabilities. Here we provide a simpler and more statistically transparent way to express the constrained space.

In Section 2 of this paper we briefly introduce conditional independence models on trees. We then proceed to describe a convenient change of coordinates for the models under consideration following [27]. In the new coordinate system the parametrization of the model has an elegant product form and in Section 3 we show how to use this to obtain the full semi-algebraic description of a simple naive Bayes model. In Section 4 we state the main result of the paper given by Theorem 4.6 and give some necessary constraints on the probability distributions in the model class using a correspondence with tree metrics. In Section 5 we discuss these results for a simple quartet tree model. Finally, in Section 6 we use the parametrization developed earlier to find an alternative form of equations given by Allman and Rhodes [1]. Our alternative specification is simpler from the algebraic point of view and has a more transparent statistical interpretation. We prove our main theorem in Section B. The paper is concluded with a short discussion.

2. TREE MODELS AND TREE CUMULANTS

In this paper we always assume that random variables are binary taking values either 0 or 1. We consider models with *hidden* variables, i.e. variables whose values are never directly observed. The vector Y has as its components all variables in the graphical model, both those that are observed and those that are hidden. The subvector of Y of observed variables is denoted by X and the subvector of hidden variables by H .

A (*directed*) *tree* $T = (V, E)$, where V is the set of vertices and $E \subseteq V \times V$ is the set of edges of T , is a connected (*directed*) graph with no cycles. A *rooted tree* is a directed tree that has one distinguished vertex called the *root*, denoted by the letter r , and all the edges are directed away from r . A rooted tree is usually denoted by T^r . By $\text{pa}(v)$ we denote the node preceding v in T^r . In particular $\text{pa}(r) = \emptyset$. A vertex of T of degree one is called a *leaf*. A vertex of T that is not a leaf is called an *inner node*.

A Markov process on a rooted tree T^r is a sequence $\{Y_v : v \in V\}$ of random variables such that for each $(\alpha_1, \dots, \alpha_{|V|}) \in \{0, 1\}^{|V|}$

$$(1) \quad p_\alpha(\theta) = \prod_{v \in V} \theta_{\alpha_v | \alpha_{\text{pa}(v)}}^{(v)},$$

where $\theta = (\theta_{\alpha_v | \alpha_{\text{pa}(v)}}^{(v)})$ and $\theta_{\alpha_v | \alpha_{\text{pa}(v)}}^{(v)} = \mathbb{P}(Y_v = \alpha_v | Y_{\text{pa}(v)} = \alpha_{\text{pa}(v)})$. Since $\theta_0^{(r)} + \theta_1^{(r)} = 1$ and $\theta_{0|i}^{(v)} + \theta_{1|i}^{(v)} = 1$ for all $v \in V \setminus \{r\}$ and $i = 0, 1$ then the set of parameters consists of exactly $2|E| + 1$ free parameters: we have two parameters: $\theta_{1|0}^{(v)}, \theta_{1|1}^{(v)}$ for each edge $(u, v) \in E$ and one parameter $\theta_1^{(r)}$ for the root. We denote the parameter space by $\Theta_T = [0, 1]^{2|E|+1}$ and the model given by this parameterization by $\widehat{\mathcal{M}}_T$.

Let n be the number of leaves of T and let $\Delta_{2^n-1} = \{p \in \mathbb{R}^{2^n} : \sum_\beta p_\beta = 1, p_\beta \geq 0\}$ with indices β ranging over $\{0, 1\}^n$ be the probability simplex of all possible distributions of $X = (X_1, \dots, X_n)$ represented by the leaves of T . Equation (1) induces a polynomial map $f_T : \Theta_T \rightarrow \Delta_{2^n-1}$ obtained by marginalization over all

the inner nodes of T


$$(2) \quad p_{\alpha_{[n]}}(\theta) = \sum_{\mathcal{H}} \prod_{v \in V} \theta_{\alpha_v | \alpha_{\text{pa}(v)}}^{(v)},$$

where $\mathcal{H}$ are all possible states of the vector of hidden variables, i.e. the sum is over $\alpha_{V \setminus [n]} \in \{0, 1\}^{|V| - n}$ and for any $A \subseteq V$, $\alpha_A = (\alpha_i)_{i \in A}$. We denote $\mathcal{M}_T = f_T(\Theta_T)$ calling it the general Markov model (c.f. [21, Section 8.3]). A *semi-algebraic set* in $\mathbb{R}^d$ is any space given by a finite number of polynomial equations and inequalities. Since Θ_T is a semi-algebraic set and f_T is a polynomial map then by the Tarski-Seidenberg theorem [3, Section 2.5.2] $\mathcal{M}_T$ is a semi-algebraic set as well.

In [27] we described a convenient change of coordinates for directed tree models as a function of the usual parametrization (2) which is expressed in terms of the probabilities. The idea was to define a regular one-to-one polynomial map $f_{p\kappa}$ from $\Delta_{2^n - 1}$ to the space of new parameters called tree cumulants $\mathcal{K}_T$. We defined a partially ordered set (poset) of all the partitions of the set of leaves induced by removing inner edges of the given tree T . Then tree cumulants are given as a one-to-one function of probabilities induced by a Möbius function on the poset. The details of this change of coordinates are given Appendix A and are illustrated below.

The tree cumulants are given by $2^n - 1$ coordinates: $\bar{\mu}_i$ for all $i \in [n]$ and a set of real-valued parameters $\{\kappa_I : I \subseteq [n] \text{ where } |I| \geq 2\}$. The first n coordinates are linear functions of the means of the n observed variables since $\bar{\mu}_i = 1 - 2\mathbb{E}X_i$. Each formula for κ_I is expressed as a function of the higher order central moments of the observed variables. These formulas are given explicitly in (31) of Appendix A. By $\mathcal{M}_T^\kappa$ we will denote the image of the original model $\mathcal{M}_T$ in the space of tree cumulants. We note that since $f_{p\kappa}$ is a one-to-one polynomial map then by the Tarski-Seidenberg theorem (see [3, Section 2.5.2]) $\mathcal{M}_T^\kappa$ is a semi-algebraic set. In this paper we provide the full semi-algebraic description of $\mathcal{M}_T^\kappa$, i.e. the complete set of polynomial equations and inequalities involving the tree cumulants which describes $\mathcal{M}_T^\kappa$ as the subset of $\mathcal{K}_T$.

Example 2.1. Consider the quartet tree model, i.e. the general Markov model given by the following graph (c.f. Section 6 in [27]).



The tree cumulants are given by 15 coordinates: $\bar{\mu}_i = 1 - 2\mathbb{P}(X_i = 1)$ for $i = 1, 2, 3, 4$ and κ_I for $I \subseteq [4]$ such that $|I| \geq 2$. Denoting $U_i = X_i - \mathbb{E}X_i$ we have $\kappa_{ij} = \mathbb{E}U_i U_j = \text{Cov}(X_i, X_j)$ for $1 \leq i < j \leq 4$ and

$$\kappa_{ijk} = \mathbb{E}(U_i U_j U_k)$$

for all $1 \leq i < j < k \leq 4$ which we note is a third order central moment. However tree cumulants of higher order cannot be equated to corresponding central moments but only expressed as functions of them. These functions are obtained by performing an appropriate Möbius inversion (see [27]). Thus for example from Appendix

A we have that

$$\kappa_{1234} = \mathbb{E}(U_1 U_2 U_3 U_4) - \mathbb{E}(U_1 U_2) \mathbb{E}(U_3 U_4).$$

But note that since the observed higher order central moments can be expressed as functions of probabilities, tree cumulants being functions of such moments can also be expressed as functions of these probabilities.

Now let Ω_T denote the set of parameters with coordinates given by $\bar{\mu}_v$ for $v \in V$ and $\eta_{u,v}$ for $(u, v) \in E$. Define a reparametrization map $f_{\theta\omega} : \Theta_T \rightarrow \Omega_T$ as follows: for every directed edge $(u, v) \in E$

$$(3) \quad \begin{aligned} \eta_{u,v} &= \theta_{1|1}^{(v)} - \theta_{1|0}^{(v)} \quad \text{and} \\ \bar{\mu}_v &= 1 - 2\lambda_v \quad \text{for each } v \in V, \end{aligned}$$

where $\lambda_v = \mathbb{E}Y_v$ is a polynomial in the original parameters θ of degree depending on the distance of v from the root r . Indeed, let $r, v_1, \dots, v_k, v$ be a directed path in T . Then

$$(4) \quad \lambda_v = \mathbb{P}(Y_v = 1) = \sum_{\alpha \in \{0,1\}^{k+1}} \theta_{1|\alpha_k}^{(v)} \theta_{\alpha_k|\alpha_{k-1}}^{(v_k)} \dots \theta_{\alpha_r}^{(r)}.$$

It can be easily checked that $\eta_{u,v} = \text{Cov}(Y_u, Y_v)/\text{Var}(Y_u)$ and hence

$$(1 - \bar{\mu}_u^2)\eta_{u,v} = (1 - \bar{\mu}_v^2)\eta_{v,u}.$$

It also follows that $\eta_{u,v}$ is just the linear regression coefficient of Y_v with respect to Y_u , namely $\mathbb{E}(Y_v - \mathbb{E}Y_v|Y_u) = \eta_{u,v}(Y_u - \mathbb{E}Y_u)$, which gives a clear statistical interpretation for the new parameters. The parameter space Ω_T is given by the following constraints: $\bar{\mu}_v \in [-1, 1]$ for all $v \in V$ and

$$(5) \quad -\min\{(1 + \bar{\mu}_u)(1 + \bar{\mu}_v), (1 - \bar{\mu}_u)(1 - \bar{\mu}_v)\} \leq (1 - \bar{\mu}_u^2)\eta_{u,v} \leq \min\{(1 + \bar{\mu}_u)(1 - \bar{\mu}_v), (1 - \bar{\mu}_u)(1 + \bar{\mu}_v)\}.$$

In [27] we proved that there is a one-to-one polynomial map between the two spaces giving the following diagram.

$$(6) \quad \begin{array}{ccc} \Theta_T & \xrightarrow{f_T} & \Delta_{2^n-1} \\ f_{\omega\theta} \uparrow & \downarrow f_{\theta\omega} & \uparrow f_{\kappa p} \downarrow f_{p\kappa} \\ \Omega_T & \xrightarrow{\psi_T} & \mathcal{K}_T \end{array}$$

One motivation behind the change of coordinates and parameters is that the induced parametrization $\psi_T : \Omega_T \rightarrow \mathcal{K}_T$ has a particularly elegant form in terms of the new parameters.

Proposition 2.2 ([27], Proposition 3). *Let $T = (V, E)$ be a rooted trivalent tree with n leaves. Then $\mathcal{M}_T^\kappa$ is parametrized by the map $\psi_T : \Omega_T \rightarrow \mathcal{K}_T$ given as an identity on the first n coordinates corresponding to $\bar{\mu}_i$ for $i \in [n]$ and on the other coordinates it is given by*

$$(7) \quad \kappa_I = \frac{1}{4} \left(1 - \bar{\mu}_{r(I)}^2\right) \prod_{v \in N(I)} \bar{\mu}_v^{\deg(v)-2} \prod_{(u,v) \in E(I)} \eta_{u,v} \quad \text{for } I \subseteq [n], |I| \geq 2$$

where the degree is taken in $T(I) = (V(I), E(I))$; $N(I)$ denotes the set of inner nodes of $T(I)$ and $r(I)$ denotes the root of $T(I)$.

Example 2.3. Consider the model in Example 2.1. The model is parametrized as in (2) by the root distribution and transition matrices attached to each of the edges. Say $\theta_1^{(r)} = 0.8$, $\theta_{1|0}^{(1)} = 0.8$, $\theta_{1|1}^{(1)} = 0.3$, $\theta_{1|0}^{(2)} = 0.7$, $\theta_{1|1}^{(2)} = 0.3$, $\theta_{1|0}^{(a)} = 0.8$, $\theta_{1|1}^{(a)} = 0.3$, $\theta_{1|0}^{(3)} = 0.7$, $\theta_{1|1}^{(3)} = 0.3$, $\theta_{1|0}^{(4)} = 0.7$, $\theta_{1|1}^{(4)} = 0.3$. Using (2) we provide the corresponding probabilities over the observed nodes in the third column in the table below. The change of coordinates presented in Appendix A gives the corresponding non-central moments $\lambda_I = \mathbb{E}(\prod_{i \in I} X_i)$ and tree cumulants κ_I supplemented with the means $\lambda_1, \lambda_2, \lambda_3, \lambda_4$.

α	I	p_α	λ_I	κ_I
0000	$\emptyset$	0.0444	1.0000	1.0000
0001	4	0.0307	0.5800	0
0010	3	0.0307	0.5800	0
0011	34	0.0403	0.3700	0.0336
0100	2	0.0346	0.6200	0
0101	24	0.0323	0.3724	0.0128
0110	23	0.0323	0.3724	0.0128
0111	234	0.0547	0.2422	-0.0020
1000	1	0.0482	0.7000	0
1001	14	0.0491	0.4220	0.0160
1010	13	0.0491	0.4220	0.0160
1011	134	0.0875	0.2750	-0.0026
1100	12	0.0828	0.4660	0.0320
1101	124	0.0979	0.2853	-0.0038
1110	123	0.0979	0.2853	-0.0038
1111	1234	0.1875	0.1875	0.0006

The formula (3) gives: $\eta_{r,1} = 0.5$, $\eta_{r,2} = 0.4$, $\eta_{r,a} = 0.5$, $\eta_{a,3} = 0.4$, $\eta_{a,4} = 0.4$ and $\bar{\mu}_1 = -0.4$, $\bar{\mu}_2 = -0.24$, $\bar{\mu}_3 = -0.16$, $\bar{\mu}_4 = -0.16$, $\bar{\mu}_r = -0.6$, $\bar{\mu}_a = -0.4$. It is easy to verify (7). For example

$$\kappa_{1234} = \frac{1}{4}(1 - \bar{\mu}_r^2)\bar{\mu}_r\bar{\mu}_a\eta_{r,1}\eta_{r,2}\eta_{r,a}\eta_{a,3}\eta_{a,4} = 0.0006.$$

Proposition 2.2 has been formulated for trivalent trees. However it can be easily extended to a more general case. For a given tree a *contraction of an edge* (u, v) results in another tree obtained from the original tree by identifying the nodes u and v and removing the edge (u, v) . Let T be a tree and let $\bar{T}$ be any trivalent tree such that T is obtained from $\bar{T}$ by edge contractions. Then $\mathcal{M}_T^\kappa \subseteq \mathcal{M}_{\bar{T}}^\kappa \subset \mathcal{K}_{\bar{T}}$ and by Corollary 4 in [27] the parameterization in (7) remains valid for T but expressed in the coordinates of $\mathcal{K}_{\bar{T}}$.

Example 2.4. Let T be a star tree with four leaves, i.e. a tree with one inner node r and four leaves connected to r by edges (r, i) for $i = 1, 2, 3, 4$. This tree can be obtained from the quartet tree in Example 2.1, denote it by $\bar{T}$, by contracting (r, a) . The model of the star tree can be realized as a subset of $\mathcal{K}_{\bar{T}}$, i.e. the space of tree cumulants for the quartet tree. The coordinates of $\mathcal{K}_{\bar{T}}$ are obtained in Example 2.1 and the parametrization of $\mathcal{M}_{\bar{T}}^\kappa$ is given for example by

$$\kappa_{1234} = \frac{1}{4}(1 - \bar{\mu}_r^2)\bar{\mu}_r^2\eta_{r,1}\eta_{r,2}\eta_{r,3}\eta_{r,4}.$$

Note however that this star tree may be obtained from many different trivalent trees by edge contraction. It follows that there exist many ways to embed the model and retain the parametrization.

Remark 2.5. Let T^r be a rooted tree and T its undirected version. Then $\mathcal{M}_T^\kappa$ depends only on T . Indeed, without loss fix two different rootings r and r' . Let T be a tree rooted in r and by T' denote its copy rooted in r' . Then $\mathcal{M}_T^\kappa = \mathcal{M}_{T'}^\kappa$, and the parameters $(\eta_{u,v}), (\bar{\mu}_v)$ and $(\eta'_{u,v}), (\bar{\mu}'_v)$ are related as follows. We have $\bar{\mu}_v = \bar{\mu}'_v$ for all $v \in V$. Moreover, if $(u, v) \in E \cap E'$ then $\eta_{u,v} = \eta'_{u,v}$ and if $(u, v) \in E \setminus E'$ then $(1 - \bar{\mu}_u^2)\eta_{u,v} = (1 - \bar{\mu}'_v{}^2)\eta'_{v,u}$. Note however that if for example $\eta_{u,v} = 0$ and $\bar{\mu}_v = 1$ then $\eta'_{v,u}$ is not well defined and in this case we set $\eta'_{v,u} = 0$. From the form of inequalities in (5) constraints on $\bar{\mu}_v$ and $\eta_{u,v}$ are satisfied if and only if the constraints on $\bar{\mu}'_v$ and $\eta'_{u,v}$ are satisfied.

3. THE SEMI-ALGEBRAIC DESCRIPTION OF THE TRIPOD TREE MODEL

In this section we obtain the full semi-algebraic description of the tripod tree model. This result is not new (see [2] [22]). However it is convenient to give a new proof of this result both to unify notation and to introduce the strategy which is used to attack the general case later. We begin with a definition.

Definition 3.1. Let A be a $2 \times 2 \times 2$ table. The hyperdeterminant of A as defined by Gelfand, Kapranov, Zelevinsky [13, Chapter 14] is given by

$$\begin{aligned} \text{Det } A &= (a_{000}^2 a_{111}^2 + a_{001}^2 a_{110}^2 + a_{010}^2 a_{101}^2 + a_{011}^2 a_{100}^2) \\ &\quad - 2(a_{000} a_{001} a_{110} a_{111} + a_{000} a_{010} a_{101} a_{111} + a_{000} a_{011} a_{100} a_{111} \\ &\quad + a_{001} a_{010} a_{101} a_{110} + a_{001} a_{011} a_{110} a_{100} + a_{010} a_{011} a_{101} a_{100}) \\ &\quad + 4(a_{000} a_{011} a_{101} a_{110} + a_{001} a_{010} a_{100} a_{111}). \end{aligned}$$

If $\sum a_{ijk} = 1$ then treating all entries formally as joint cell probabilities (without positivity constraints) we can simplify this formula using the change of coordinates to central moments. The reparameterizations in Appendix A are well defined for this extended space of probabilities and we have that

$$(8) \quad \text{Det } A = \mu_{123}^2 + 4\mu_{12}\mu_{13}\mu_{23},$$

which can be verified by direct computations. We note in passing that a similar idea of treating moments formally lies behind the umbral calculus [20].

From the construction of tree cumulants (c.f. Appendix A) it follows that $\kappa_I = \mu_I$ for all $I \subseteq [n]$ such that $|I| \leq 3$. Henceforth, for clarity, these lower order tree cumulants will be written as their more familiar corresponding central moments.

Lemma 3.2 (The semi-algebraic description of the tripod model). *Let $\mathcal{M}_T$ be the general Markov model on a tripod tree T rooted in any node of T . Let P be a $2 \times 2 \times 2$ probability table for three binary random variables (X_1, X_2, X_3) with central moments $\mu_{12}, \mu_{13}, \mu_{23}, \mu_{123}$ (equivalent to the corresponding tree cumulants) and $\bar{\mu}_i = 1 - 2\mathbb{E}X_i$ for $i = 1, 2, 3$. Then $\mathcal{M}_T^\kappa$ is given by*

$$(9) \quad \begin{aligned} \mu_{ij} &= \frac{1}{4}(1 - \bar{\mu}_h^2)\eta_{h,i}\eta_{h,j} \text{ for all } i \neq j \in \{1, 2, 3\} \text{ and} \\ \mu_{123} &= \frac{1}{4}(1 - \bar{\mu}_h^2)\bar{\mu}_h\eta_{h,1}\eta_{h,2}\eta_{h,3}, \end{aligned}$$

where $\bar{\mu}_h \in [-1, 1]$ and $(1 - \bar{\mu}_h^2)\eta_{h,i}$ for all $i = 1, 2, 3$ satisfy the inequality (5). Moreover, $P \in \mathcal{M}_T$ if and only if $K = f_{p\kappa}(P) \in \mathcal{K}_T = \mathcal{C}_3$ satisfies the following inequalities

$$(10) \quad \mu_{12}\mu_{13}\mu_{23} \geq 0,$$

$$(11) \quad \mu_{12}^2\mu_{13}^2 + \mu_{12}^2\mu_{23}^2 + \mu_{13}^2\mu_{23}^2 \leq \text{Det } P \leq \min_{1 \leq i < j \leq 3} \mu_{ij}^2$$

and

$$(12) \quad \text{Det } P \leq \min \left\{ ((1 + \bar{\mu}_i)\mu_{jk} - \mu_{123})^2, ((1 - \bar{\mu}_i)\mu_{jk} + \mu_{123})^2 \right\},$$

for all $i = 1, 2, 3$ where by j, k we denote elements of $\{1, 2, 3\} \setminus i$.

Proof. By Remark 2.5 $\mathcal{M}_T^\kappa$ does not depend on the rooting and hence we can assume that T is rooted in h . The parameterization in (9) follows from Proposition 2.2 by considering T rooted at h and the corresponding independence statements $X_1 \perp\!\!\!\perp (X_2, X_3) | H$ and $X_2 \perp\!\!\!\perp X_3 | H$.

Denote by $\mathcal{M}$ the subset of $\mathcal{K}_T$ given by inequalities in (10), (11) and (12). We need to show that $\mathcal{M} = \mathcal{M}_T^\kappa$ for any rooting of T . First we prove that $\mathcal{M}_T^\kappa \subseteq \mathcal{M}$. Let $K = \psi_T(\omega)$ for some $\omega \in \Omega_T$ with coordinates given by $\bar{\mu}_h$ and $\bar{\mu}_i, \eta_{h,i}$ for $i = 1, 2, 3$. Using (9) we obtain

$$(13) \quad \mu_{12}\mu_{13}\mu_{23} = \left(\frac{1}{4}(1 - \bar{\mu}_h^2) \right)^3 (\eta_{h,1}\eta_{h,2}\eta_{h,3})^2.$$

Since $\bar{\mu}_h \in [-1, 1]$ this implies the inequality in (10). Moreover, we have

$$(14) \quad \text{Det } P = \mu_{123}^2 + 4\mu_{12}\mu_{13}\mu_{23} = \frac{1}{16}(1 - \bar{\mu}_h^2)^2(\eta_{h,1}\eta_{h,2}\eta_{h,3})^2.$$

Multiplying both sides by $\bar{\mu}_h^2$ together with the second equation in (9) implies

$$(15) \quad \bar{\mu}_h^2 \text{Det } P = \mu_{123}^2, \quad (1 - \bar{\mu}_h^2) \text{Det } P = 4\mu_{12}\mu_{13}\mu_{23}.$$

On the other hand (9) and (14) imply also that

$$(16) \quad \eta_{h,i}^2 \mu_{jk}^2 = \text{Det } P \quad \text{for all } i = 1, 2, 3.$$

Again by substituting μ_{ij} for $\frac{1}{4}(1 - \bar{\mu}_h^2)\eta_{h,i}\eta_{h,j}$ it can be shown that

$$(17) \quad \mu_{12}^2\mu_{13}^2 + \mu_{12}^2\mu_{23}^2 + \mu_{13}^2\mu_{23}^2 = \frac{1}{16}(1 - \bar{\mu}_h^2)^2(\eta_{h,1}^2 + \eta_{h,2}^2 + \eta_{h,3}^2)\text{Det } P.$$

Since necessarily $\eta_{h,i}^2, \bar{\mu}_h^2 \in [0, 1]$ then (15), (16) and (17) imply that

$$\mu_{12}^2\mu_{13}^2 + \mu_{12}^2\mu_{23}^2 + \mu_{13}^2\mu_{23}^2 \leq \text{Det } P \leq \min_{i,j} \mu_{ij}^2.$$

To show that K satisfies (12) first divide this inequality by μ_{jk}^2 (if it zero the inequality is trivially satisfied since $\text{Det } P = \mu_{123}^2$). Using (16) and the fact that $\frac{\mu_{123}}{\mu_{jk}} = \bar{\mu}_h\eta_{h,i}$ we obtain

$$\eta_{h,i}^2 \leq \min\{((1 + \bar{\mu}_i) - \bar{\mu}_h\eta_{h,i})^2, ((1 - \bar{\mu}_i) + \bar{\mu}_h\eta_{h,i})^2\}.$$

This is equivalent to

$$\begin{aligned} (\eta_{h,i}(1 - \bar{\mu}_h^2) + \bar{\mu}_h(1 + \bar{\mu}_i))^2 &\leq (1 + \bar{\mu}_i)^2, \\ (\eta_{h,i}(1 - \bar{\mu}_h^2) - \bar{\mu}_h(1 - \bar{\mu}_i))^2 &\leq (1 - \bar{\mu}_i)^2. \end{aligned}$$

Since $1 \pm \bar{\mu}_i \geq 0$ this in turn reduces to (5). Therefore since by hypothesis (5) holds and hence $\mathcal{M}_T^\kappa \subseteq \mathcal{M}$.

Now we show that $\mathcal{M} \subseteq \mathcal{M}_T^\kappa$ by proving that for $K \in \mathcal{M}$ a parameter ω in (9) exists which satisfies constraints defining Ω_T and $K = \psi_T(\omega)$. Let $P = f_{p^\kappa}^{-1}(K)$ then from (10) we know that $\text{Det } P \geq 0$. So consider separately the two situations: first when $\text{Det } P = 0$ and second when $\text{Det } P > 0$. In the first case again from (10) necessarily $\mu_{123} = 0$. The inequality (11) therefore implies that at least two covariances are zero. If all the covariances are zero then for $\eta_{h,1} = \eta_{h,2} = \eta_{h,3} = 0$ and $\bar{\mu}_h^2 = 1$ we obtain a valid choice of parameters in (9) and the values satisfy (5). When one covariance, say $\mu_{12} \neq 0$, is non-zero then if a choice of parameters exists it has to satisfy $\bar{\mu}_h^2 \neq 1$, $\eta_{h,1}, \eta_{h,2} \neq 0$ and $\eta_{h,3} = 0$. Such a choice of parameters will exist if we can ensure that $\mu_{12} = (1 - \bar{\mu}_h^2)\eta_{h,1}\eta_{h,2}$. This follows from Corollary 2 in [14] which states that if only $\mu_{12} \neq 0$ then there always exists a choice of parameters for model $X_1 \perp\!\!\!\perp X_2 | H$, where H is hidden.

Assume now that $\text{Det } P > 0$ which by (11) implies that $\mu_{ij} \neq 0$ for each $i < j = 1, 2, 3$. Set $\bar{\mu}_h^2 = \frac{\mu_{123}^2}{\text{Det } P}$ and $\eta_{h,i}^2 = \frac{\text{Det } P}{\mu_{jk}^2}$ for $i = 1, 2, 3$. It follows that $(\frac{1}{4}(1 - \bar{\mu}_h^2))^2 \eta_{h,i}^2 \eta_{h,j}^2 = \mu_{ij}^2$ for $i, j = 1, 2, 3$ and $(\frac{1}{4}(1 - \bar{\mu}_h^2))^2 \bar{\mu}_h^2 \eta_{h,1}^2 \eta_{h,2}^2 \eta_{h,3}^2 = \mu_{123}^2$, which coincides with (9) modulo the sign. It can be easily shown that $\mu_{12}\mu_{13}\mu_{23} > 0$ implies that there exist a choice of signs for $\eta_{h,i}$ for $i = 1, 2, 3$ such that

$$\frac{1}{4}(1 - \bar{\mu}_h^2)\eta_{h,i}\eta_{h,j} = \mu_{ij}$$

for all $1 \leq i < j \leq 3$ as in (9). For example set $\text{sgn}(\eta_{h,i}) = \text{sgn}(\mu_{jk})$ and use the fact that by our assumption $\text{sgn}(\mu_{ij}) = \text{sgn}(\mu_{ik})\text{sgn}(\mu_{jk})$. This choice of signs already determines the sign of $\bar{\mu}_h$ so that

$$\frac{1}{4}(1 - \bar{\mu}_h^2)\bar{\mu}_h\eta_{h,1}\eta_{h,2}\eta_{h,3} = \mu_{123}$$

holds.

It remains to show that parameters set in this way satisfy the constraints defining Ω_T . First note that since $0 \leq 4\mu_{12}\mu_{13}\mu_{23} \leq \text{Det } P$ then $\bar{\mu}_h^2 \in [0, 1]$ as required. From Appendix D in [27] we know that if $(\eta_{h,1}, \eta_{h,2}, \eta_{h,3}, \bar{\mu}_h)$ is one choice of parameters then there exists only one alternative choice and it is $(-\eta_{h,1}, -\eta_{h,2}, -\eta_{h,3}, -\bar{\mu}_h)$. For each $i = 1, 2, 3$ we check if $(1 - \bar{\mu}_h^2)\eta_{h,i}$ satisfies (5). For a fixed $i = 1, 2, 3$ one easily checks that $(\eta_{h,i}, \bar{\mu}_h)$ satisfies (5) if and only if $(-\eta_{h,i}, -\bar{\mu}_h)$ does. Therefore one can assume that $\eta_{h,i} = \frac{\sqrt{\text{Det } P}}{|\mu_{jk}|} > 0$. In this case $\bar{\mu}_h = s(j, k) \frac{\mu_{123}}{\sqrt{\text{Det } P}}$ where $s(j, k) = \text{sgn}(\mu_{jk})$. From this it follows that (5) is satisfied if and only if

$$(18) \quad 4\mu_{12}\mu_{13}\mu_{23} \leq (1 \pm \bar{\mu}_i)(\sqrt{\mu_{jk}^2 \text{Det } P} \mp \mu_{jk}\mu_{123}).$$

We show that (10) and (12) already imply that this inequality has to be satisfied. Multiply both sides by $\sqrt{\mu_{jk}^2 \text{Det } P} \pm \mu_{jk}\mu_{123}$ and then divide by $4\mu_{12}\mu_{13}\mu_{23}$ (both expressions are strictly positive) to obtain

$$(19) \quad \sqrt{\mu_{jk}^2 \text{Det } P} \leq (1 \pm \bar{\mu}_i)\mu_{jk}^2 \mp \mu_{jk}\mu_{123}.$$

Since $\text{Det}P > 0$ and $\mu_{jk}^2 > 0$ this is equivalent to

$$(20) \quad \begin{aligned} 0 &\leq (1 \pm \bar{\mu}_i)\mu_{jk}^2 \mp \mu_{jk}\mu_{123}, \\ \text{Det}P &\leq ((1 \pm \bar{\mu}_i)\mu_{jk} \mp \mu_{123})^2 \end{aligned}$$

The second inequality in (20) is exactly (12). So we have to show that the first inequality is already implied by (10) and (12). To see this rewrite (12) as

$$4\mu_{12}\mu_{13}\mu_{23} \leq (1 \pm \bar{\mu}_i)^2\mu_{jk}^2 \mp 2\mu_{jk}\mu_{123}(1 \pm \bar{\mu}_i).$$

However, since $\mu_{12}\mu_{13}\mu_{23} \geq 0$ the right hand side of the inequality above has to be non-negative as well. In particular

$$(21) \quad \begin{aligned} (1 - \bar{\mu}_i)\mu_{jk}^2 &\geq -2\mu_{123}\mu_{jk} \\ (1 + \bar{\mu}_i)\mu_{jk}^2 &\geq 2\mu_{123}\mu_{jk}. \end{aligned}$$

noting that the left-hand sides are nonnegative. For each of the two inequalities if the right-hand side is negative then the inequality is trivially satisfied. If the right-hand side is nonnegative then in the first case $-2\mu_{123}\mu_{jk} \geq -\mu_{123}\mu_{jk}$ and in the second case $2\mu_{123}\mu_{jk} \geq \mu_{123}\mu_{jk}$. Hence the following set of inequalities is implied by (21)

$$(1 - \bar{\mu}_i)\mu_{jk}^2 \geq -\mu_{123}\mu_{jk}, \quad (1 + \bar{\mu}_i)\mu_{jk}^2 \geq \mu_{123}\mu_{jk}.$$

This is exactly the first inequality in (20) which shows that it is implied by (10) and (12). Consequently, (18) and hence also (5) are satisfied. It follows that $\mathcal{M} \subseteq \mathcal{M}_T^\kappa$. $\square$

4. A CONNECTION WITH TREE METRICS

Now let T be a general tree with n leaves. Before stating the main theorem of the paper we first show how to obtain an elegant set of necessary constraints on $\mathcal{M}_T$. In this section we assume that $\bar{\mu}_v^2 \neq 1$ for all $v \in V$ (c.f. Remark 4.7). Since $\text{Var}(Y_u) = \frac{1}{4}(1 - \bar{\mu}_u^2)$ the correlation between Y_u and Y_v is defined as $\rho_{uv} = \frac{4\mu_{uv}}{\sqrt{(1 - \bar{\mu}_u^2)(1 - \bar{\mu}_v^2)}}$ which gives

$$(22) \quad \rho_{uv} = \eta_{u,v} \sqrt{\frac{1 - \bar{\mu}_u^2}{1 - \bar{\mu}_v^2}} = \eta_{v,u} \sqrt{\frac{1 - \bar{\mu}_v^2}{1 - \bar{\mu}_u^2}}.$$

Lemma 4.1. *For any $i, j \in [n]$ let $E(ij)$ be the set of edges on the unique path joining i and j in T . Then*

$$(23) \quad \rho_{ij} = \prod_{(u,v) \in E(ij)} \rho_{uv}$$

for each probability distribution in $\mathcal{M}_T^\kappa$ such that all the correlations are well defined.

Proof. By (7) applied to $T(ij)$ we have $\mu_{ij} = \frac{1}{4}(1 - \bar{\mu}_r^2) \prod_{(u,v) \in E(ij)} \eta_{u,v}$, where r is the root of the path between i and j and hence

$$\rho_{ij} = \sqrt{\frac{1 - \bar{\mu}_r^2}{1 - \bar{\mu}_i^2}} \sqrt{\frac{1 - \bar{\mu}_r^2}{1 - \bar{\mu}_j^2}} \prod_{(u,v) \in E(ij)} \eta_{u,v}.$$

Now apply (22) to each $\eta_{u,v}$ in the product above to show (23). $\square$

The above equation allows us to demonstrate an interesting reformulation of our problem in term of tree metrics (c.f. [21, Section 7]) which we explain below (see also Cavender [7]).

Definition 4.2. An arbitrary function $\delta : [n] \times [n] \rightarrow \mathbb{R}$ is called a tree metric if there exists a tree $T = (V, E)$ with the set of leaves given by $[n]$ and with a positive real-valued weighting $w : E \rightarrow \mathbb{R}_{>0}$ such that for all $i, j \in [n]$

$$\delta(i, j) = \begin{cases} \sum_{e \in E(ij)} w(e), & \text{if } i \neq j, \\ 0, & \text{otherwise} \end{cases}$$

Let $d : V \times V \rightarrow \mathbb{R}$ be a map defined as

$$d(k, l) = \begin{cases} -\log(\rho_{kl}^2), & \text{for all } k, l \in V \text{ such that } \rho_{kl} \neq 0, \\ +\infty, & \text{otherwise} \end{cases}$$

then $d(k, l) \geq 0$ because $\rho_{kl}^2 \leq 1$ and $d(k, k) = 0$ for all $k \in V$ since $\rho_{kk} = 1$. If $K \in \mathcal{M}_T^\kappa$ then by (23) $\rho_{ij}^2 = \prod_{e \in E(ij)} \rho_e^2$ and we can define map $d_{(T;K)} : [n] \times [n] \rightarrow \mathbb{R}$

$$(24) \quad -\log(\rho_{ij}^2) = d_{(T;K)}(i, j) = \begin{cases} \sum_{(u,v) \in E(ij)} d(u, v), & \text{if } i \neq j, \\ 0, & \text{otherwise.} \end{cases}$$

This map is a tree metric by Definition 4.2. In our case we have a point in the model space defining all the second order correlations and $d_{(T;K)}(i, j)$ for $i, j \in [n]$. The question is: What are the conditions for the “distances” between leaves so that there exists a tree T and edge lengths $d(u, v)$ for all $(u, v) \in E$ such that (24) is satisfied? Or equivalently: What are the conditions on the absolute values of the second order correlations in order that $\rho_{ij}^2 = \prod_{e \in E_{ij}} \rho_e^2$ (for some edge correlations) is satisfied? We have the following theorem.

Theorem 4.3 (Tree-Metric Theorem, Buneman [4]). *A function $\delta : [n] \times [n] \rightarrow \mathbb{R}$ is a tree metric on $[n]$ if and only if for every four (not necessarily distinct) elements $i, j, k, l \in [n]$,*

$$\delta(i, j) + \delta(k, l) \leq \max \{ \delta(i, k) + \delta(j, l), \delta(i, l) + \delta(j, k) \}.$$

Moreover, a tree metric defines the tree uniquely.

The question we may now ask is - For a given assignments of edge weights on a tree metric, which of these correspond to a probability model on the tree defined as the image of (2)? From Lemma 4.1 we have seen that the tree metric itself induces (using Definition 4.2) some necessary conditions related to the four-point condition. Since $\delta(i, j) = \log(-\rho_{ij})$ these constraints in terms of correlations translate in

$$-\log(\rho_{ij}^2 \rho_{kl}^2) \leq -\min \{ \log(\rho_{ik}^2 \rho_{jl}^2), \log(\rho_{il}^2 \rho_{jk}^2) \}.$$

Since log is a monotone function we obtain

$$(25) \quad \min \left\{ \frac{\rho_{ik}^2 \rho_{jl}^2}{\rho_{ij}^2 \rho_{kl}^2}, \frac{\rho_{il}^2 \rho_{jk}^2}{\rho_{ij}^2 \rho_{kl}^2} \right\} = \min \left\{ \frac{\mu_{ik}^2 \mu_{jl}^2}{\mu_{ij}^2 \mu_{kl}^2}, \frac{\mu_{il}^2 \mu_{jk}^2}{\mu_{ij}^2 \mu_{kl}^2} \right\} \leq 1$$

for all not necessarily distinct leaves $i, j, k, l \in [n]$. However later in Theorem 4.6 we show that these constraints are not the only active constraints on the model $\mathcal{M}_T$.

Before we present this theorem it is helpful to make some simple observations about the relationship between correlations and probabilistic tree models. Since $\rho_{u,v}$ can have different signs we define a signed tree metric as a tree metric with

an additional sign assignment for each edge of T . There are additional natural constraints which assure that there exists a choice of signs for edge correlations such that (23) is satisfied.

Lemma 4.4. *Let T be a tree with n leaves and let $s : E \rightarrow \{-1, 1\}$ be the map assigning signs to edges of T . Suppose that we have a map $\sigma : [n] \times [n] \rightarrow \{-1, 1\}$. Then there exists a map s as defined above such that for all $i, j \in [n]$*

$$(26) \quad \sigma(i, j) = \prod_{(u,v) \in E(ij)} s(u, v)$$

if and only if for all triples $i, j, k \in [n]$ $\sigma(i, j)\sigma(i, k)\sigma(j, k) = 1$.

Proof. First assume that there exists the map s as in the statement. It induces a map $s : V \times V \rightarrow \{-1, 1\}$ (we use the same notation) such that $s(k, l) = \prod_{(u,v) \in E(kl)} s(u, v)$. For any triple i, j, k there exists a unique inner node h which is the intersection of all three paths between i, j, k . By the above equation the choice of signs for all $s(u, v)$ ($u, v \in E$) gives $s(i, h), s(j, h)$ and $s(k, h)$. Since $s(i, j) = s(i, h)s(j, h)$ and the same for the two other pairs, we get that $s(i, j)s(i, k)s(j, k) = s^2(i, h)s^2(j, h)s^2(k, h) = 1$ and the result follows since by construction $\sigma(i, j) = s(i, j)$ for all $i, j \in [n]$.

To prove the converse we use an inductive argument with respect to number of hidden nodes. Note that whenever there is a path $E(uv)$ in T such that all its inner nodes have degree two then a sign assignment satisfying (26) exists if and only if there exists a sign assignment for the same tree but with $E(uv)$ contracted to a single edge (u, v) . Hence we can assume that the degree of each inner node is at least three. First we will show that the theorem is true for trees with one inner node (star trees). In this case we will use induction with respect to number of leaves. The theorem is true for the tripod tree what can be checked directly. Assume it works for all star trees with $k \leq m - 1$ leaves and let T be a star tree with m leaves. By assumption for any three leaves i, j, k : $\sigma(i, j)\sigma(i, k)\sigma(j, k) = 1$. If we consider a subtree with $(1, h)$ deleted then by induction assumption we can find a consistent choice of signs for all remaining edge correlations. A choice of a sign for $(1, h)$ consistent with (26) exists if for all $i \geq 2$ $\sigma(1, i) = s(1, h)s(i, h)$. This is true if either $\sigma(1, i)s(i, h) = 1$ for all i or $\sigma(1, i)s(i, h) = -1$ for all i . Assume it is not true, i.e. there exist two leaves i, j such that $\sigma(1, i)s(i, h) = 1$ and $\sigma(1, j)s(j, h) = -1$. Then in particular since $\sigma(i, j) = s(i, h)s(j, h)$ we have $\sigma(1, i)\sigma(1, j)\sigma(i, j) = -1$ which contradicts our assumption.

If the number of the inner nodes is greater than one, pick an inner node h adjacent to exactly one inner node. Let h' be the inner node adjacent to h and let I be a subset of leaves which are adjacent to h . Let $1 \in I$ and consider a subtree T' obtained by removing all leaves in I and the incident edges apart from 1 and $(h, 1)$. By the induction, since h has degree two in the resulting subtree, we can find signs for all edge correlations of T' . Set $s(h, h') = 1$ then $s(h, 1) = s(h', 1)$ and we need only to show that we can identify $s(h, i)$ for all $i \in I \setminus 1$. Let j, k be any two leaves not in I and let $i \in I$. Using exactly the same argument we used for the analysis of the star tree case we can now show by contradiction that for each $i \in I$, there exists an assignment of $s(i, h)s(h, h') = s(i, h)$. $\square$

The lemma implies that for all $i, j, k \in [n]$ necessarily $\rho_{ij}\rho_{ik}\rho_{jk} \geq 0$ or equivalently that for all $i, j, k \in [n]$ necessarily $\mu_{ij}\mu_{ik}\mu_{jk} \geq 0$. This in particular implies

that $\frac{\mu_{ik}\mu_{jl}}{\mu_{ij}\mu_{kl}} \geq 0$ for all $i, j, k, l \in [n]$. By taking the square root in (25) these constraints can be combined and rearranged to give the inequalities

$$(27) \quad 0 \leq \min \left\{ \frac{\mu_{ik}\mu_{jl}}{\mu_{ij}\mu_{kl}}, \frac{\mu_{il}\mu_{jk}}{\mu_{ij}\mu_{kl}} \right\} \leq 1$$

for all (not necessarily distinct) $i, j, k, l \in [n]$. We have now obtained the complete set of inequality constraints on $\mathcal{M}_T$ that involve only second order moments in their expression. However the fact that additional constraints involving higher order moments exist is illustrated in the following simple example.

Example 4.5. Consider the tripod tree model in Lemma 3.2. Let K be a point in $\mathcal{K}_T$ given by $\bar{\mu}_i = 0.7$ for $i = 1, 2, 3$, $\mu_{ij} = 0.0625$ (or equivalently $\rho_{ij} = 0.49$) for each $i < j$ and $\mu_{123} = 0.0526$. This point lies in the space of tree cumulants $\mathcal{K}_T$ which can be checked by mapping back the central moments to probabilities, since the resulting vector $[p_\alpha]$ lies in Δ_7 .

Clearly K satisfies all the tree metric constraints in (27). The equation (4.1) is satisfied with $\rho_{hi} = 0.7$ for each $i = 1, 2, 3$. We now show that despite this $K \notin \mathcal{M}_T^\kappa$. For if $K \in \mathcal{M}_T^\kappa$ we could find $\bar{\mu}_h$ and $\eta_{h,i}$ satisfying constraints in (5) so that (9) held. Using the formulas in Corollary 11 in [27] it is easy to compute that $\bar{\mu}_h = 0.86$ and $\eta_{h,i} \approx 0.98$. To confirm this substitute these values into (9) to check the equations are satisfied. However, K is not in the model since these parameters do not lie in Ω_T . Indeed,

$$(1 - \bar{\mu}_h^2)\eta_{h,i} \approx 0.255 > (1 + \bar{\mu}_i)(1 - \bar{\mu}_h) = 0.238$$

and hence (5) is not satisfied.

The consequence of the fact that the parameters do not lie in Ω_T is that this parametrization does not lead to a valid assignment of conditional probabilities to the edges of the tree. For example with numbers given above we can calculate that the induced marginal distribution for (X_i, H) would have to satisfy $\mathbb{P}(X_i = 0, H = 1) = -0.0043$ which is obviously not a consistent assignment for a probability model. Thus there must exist other constraints involving observed higher order moments that need to hold for a probability model to be valid. We note that for the tripod tree these were given by Lemma 3.2.

In Section B we prove the following theorem which gives the complete set of constraints which have to be satisfied by tree cumulants to lie in $\mathcal{M}_T$ in the case when T is a trivalent tree. For the statement of this result we need the following definitions. For each edge e of T we define the edge split $(A)(B)$ as a partition of the set of leaves into two subsets A and B which correspond to two connected components of T obtained after removing e . For example in Example 2.1 removing (r, a) induces (12)(34). Moreover, let $P \in \Delta_{2^n-1}$ be the probability distribution of the vector $(X_1, \dots, X_n)$ then for any $i, j, k \in [n]$ P^{ijk} denotes the $2 \times 2 \times 2$ table of the marginal distribution of (X_i, X_j, X_k) .

Theorem 4.6. *Let $T = (V, E)$ be a trivalent tree with n leaves. Let $\mathcal{M}_T \subseteq \Delta_{2^n-1}$ be the model defined as an image of the parametrization in (2) and $\mathcal{M}_T^\kappa = f_{p_\kappa}(\mathcal{M}_T)$. Suppose P is a joint probability distribution on n binary variables and $K = f_{p_\kappa}(P)$. Then $K \in \mathcal{M}_T^\kappa$ (or equivalently $P \in \mathcal{M}_T$) if and only if the following four conditions hold:*

(C1): For each edge split $(A)(B)$ whenever we have four nonempty subsets (not necessarily disjoint) $I_1, I_2 \subseteq A$, $J_1, J_2 \subseteq B$ then we must have that

$$\kappa_{I_1 J_1} \kappa_{I_2 J_2} - \kappa_{I_1 J_2} \kappa_{I_2 J_1} = 0.$$

(C2): For all $1 \leq i < j < k \leq n$ we have

$$\mu_{ij} \mu_{ik} \mu_{jk} \geq 0$$

and

$$(\mu_{ij}^2 \mu_{ik}^2 + \mu_{ij}^2 \mu_{jk}^2 + \mu_{ik}^2 \mu_{jk}^2) \leq \text{Det } P^{ijk} \leq \min_{l, l' \in \{i, j, k\}} \mu_{ll'}^2,$$

(C3): for all $1 \leq i < j < k \leq n$

$$\text{Det } P^{ijk} \leq ((1 \pm \bar{\mu}_{\sigma(i)}) \mu_{\sigma(j)\sigma(k)} \mp \mu_{ijk})^2,$$

for all three permutations σ of $\{i, j, k\}$ such that $\sigma(j) < \sigma(k)$.

(C4): for all $I \subseteq [n]$ if there exist $i, j \in I$ such that $\mu_{ij} = 0$ then $\kappa_I = 0$

(C5): for any $i, j, k, l \in [n]$ such that there exists $e \in E$ inducing a split $(A)(B)$ such that $i, j \in A$ and $k, l \in B$ we have

$$(2\mu_{ik}\mu_{jl})^2 \leq (\sqrt{\mu_{jl}^2 \text{Det } P^{ijk}} \pm \mu_{jl}\mu_{ijk})(\sqrt{\text{Det } P^{ikl}} \mp \mu_{ikl}).$$

Remark 4.7. In the phylogenetic analysis it is often assumed that $\eta_{u,v} > 0$ for all $(u, v) \in E$ and $\bar{\mu}_v^2 \neq 1$ for all $v \in V$ (c.f. assumptions (M1)-(M3) in Section 8.2 and Section 8.4 in [21]). In this case $\mu_{ij}\mu_{ik}\mu_{jk} > 0$ for all $i, j, k \in [n]$ and the second constraint in (C2) is not active. Moreover, the model is globally identified (c.f. Appendix D in [27]).

5. EXAMPLE: THE QUARTET TREE MODEL

We can check that modulo the numerical error $K \in \mathcal{K}_T$ provided in the table from Example 2.3 satisfies all the constraints in Theorem 4.6. To check (C1) note for example that

$$\kappa_{13}\kappa_{24} - \kappa_{14}\kappa_{23} = 0.0160 \cdot 0.0128 - 0.0160 \cdot 0.0128 = 0,$$

$$\kappa_{123}\kappa_{134} - \kappa_{1234}\kappa_{13} = (-0.00384) \cdot (-0.00256) - 0.0006144 \cdot 0.016 = 1.6941 \cdot 10^{-21} \approx 0.$$

The last equation shows that due to the limited precision of numerical software typically the equations in (C1) will not be satisfied exactly.

To check (C2) check for example that $\text{Det } P^{123} \approx 4.096 \cdot 10^{-5}$, $\min\{\mu_{12}^2, \mu_{13}^2, \mu_{23}^2\} \approx 1.6384 \cdot 10^{-4}$ and $\mu_{12}^2 \mu_{13}^2 + \mu_{12}^2 \mu_{23}^2 + \mu_{13}^2 \mu_{23}^2 \approx 4.7186 \cdot 10^{-7}$. For (C3) again we check only one of all the constraints. One has

$$((1 \pm \bar{\mu}_1) \mu_{23} \mp \mu_{123})^2 = \{1.3271 \cdot 10^{-4}, 1.9825 \cdot 10^{-4}\}$$

$$((1 \pm \bar{\mu}_2) \mu_{13} \mp \mu_{123})^2 = 2.5600 \cdot 10^{-4}$$

$$((1 \pm \bar{\mu}_3) \mu_{12} \mp \mu_{123})^2 = \{9.4372 \cdot 10^{-4}, 0.0011\}$$

and hence

$$\text{Det } P^{123} \approx 4.096 \cdot 10^{-5} \leq \min \left\{ ((1 \pm \bar{\mu}_{\sigma(i)}) \mu_{\sigma(j)\sigma(k)} \mp \mu_{ijk})^2 \right\} \approx 1.3271 \cdot 10^{-4}$$

is satisfied. In a similar way we check (C5).

If we have only tree cumulants we can still identify the parameters of the model up to the label switching on the inner nodes using Corollary 11 in [27]. For example

$$\begin{aligned}\bar{\mu}_r^2 &= \frac{\mu_{123}^2}{\mu_{123}^2 + 4\mu_{12}\mu_{13}\mu_{23}} = 0.36 \\ \eta_{r,1}^2 &= \frac{\mu_{123}^2 + 4\mu_{12}\mu_{13}\mu_{23}}{\mu_{23}^2} = 0.25 \\ \eta_{r,a}^2 &= \frac{\mu_{14}^2 \mu_{123}^2 + 4\mu_{12}\mu_{13}\mu_{23}}{\mu_{12}^2 \mu_{134}^2 + 4\mu_{13}\mu_{14}\mu_{34}} = 0.25.\end{aligned}$$

Note that the numbers on the right hand side of the three equations above do not depend on the choices made. For example by Corollary 11 [27] to compute $\bar{\mu}_r^2$ we can use any three leaves separated by r . In the formula above we used 1, 2, 3 but we could also use 1, 2, 4. But in both cases we get the same result, namely that

$$\bar{\mu}_r^2 = \frac{\mu_{124}^2}{\mu_{124}^2 + 4\mu_{12}\mu_{14}\mu_{24}} = 0.36.$$

If (C1) is not satisfied but (C2)-(C5) hold then the numbers depend on the choices of leaves. However the solutions for equations can be obtained from Corollary 11 [27] and any of these choices gives feasible values for parameters in Ω_T .

From the point of view of the original motivation behind the paper a different scenario is of an interest. Imagine that we have $K \in \mathcal{K}_T$ such that all the equations in (C1) are satisfied, i.e. all the phylogenetic invariants hold. If one of the constraints in (C2)-(C5) then $K \notin \mathcal{M}_T^\kappa$. This shows that the method of phylogenetic invariants as it is commonly used may lead to spurious results. For example consider sample proportions and the corresponding tree cumulants as in the table below

α	I	p_α	λ_I	κ_I
0000	$\emptyset$	0.0755	1.0000	1.0000
0001	4	0.0483	0.5800	0
0010	3	0.0483	0.5800	0
0011	34	0.0579	0.3700	0.0336
0100	2	0.0479	0.6200	0
0101	24	0.0399	0.3724	0.0128
0110	23	0.0399	0.3724	0.0128
0111	234	0.0623	0.2422	-0.0020
1000	1	0.0171	0.5800	0
1001	14	0.0315	0.3716	0.0352
1010	13	0.0315	0.3716	0.0352
1011	134	0.0699	0.2498	-0.0056
1100	12	0.0695	0.4300	0.0704
1101	124	0.0903	0.2702	-0.0084
1110	123	0.0903	0.2702	-0.0084
1111	1234	0.1799	0.1799	0.0014

It can be checked that for this point all the equations in (C1) are satisfied. However it is not in the model. Using the formulas in Corollary 11 [27] it is simple to confirm that the point mapping to K satisfies $\theta_{1|0}^{(1)} = -0.3$. This cannot therefore be a probability and so $\theta \notin \Theta_T$.

6. PHYLOGENETIC INVARIANTS

In a seminal paper Allman and Rhodes [1] identified equations defining the general Markov $\mathcal{M}_T$ in the case when T is a trivalent tree. In this section we relate their results to ours. To introduce their main theorem we need the following definition.

Definition 6.1. Let $X = (X_1, \dots, X_n)$ be a vector of binary random variables and let $P = (p_\gamma)_{\gamma \in \{0,1\}^n}$ be a $2 \times \dots \times 2$ table of the joint distribution of X . Let $(A)(B)$ form a partition of $[n]$. Then the *flattening* of P induced by the partition is a matrix

$$P_{(A)(B)} = [p_{\alpha\beta}], \quad \alpha \in \{0,1\}^{|A|}, \beta \in \{0,1\}^{|B|},$$

where $p_{\alpha\beta} = \mathbb{P}(X_A = \alpha, X_B = \beta)$. Let $T = (V, E)$ be a tree. In particular, for each $e \in E$, removing edge e from E induces a partition of the set of leaves into two subsets corresponding to the two connected components of the resulting forest. They called this flattening an *edge flattening* and we denote it by P_e .

Note that whenever we implicitly use some order on coordinates indexed by $\{0,1\}$ -sequences we always mean the order induced by the lexicographic order on $\{0,1\}$ -sequences such that $0 \dots 00 > 0 \dots 01 > \dots > 1 \dots 11$.

If P is the joint distribution of $X = (X_1, \dots, X_n)$ then each of its flattenings is just a matrix representation of the joint distribution P and contains essentially the same probabilistic data. However, these different representations contain important geometric information about the model.

Theorem 6.2 (Allman, Rhodes [1]). *Let T^r be a trivalent tree rooted in r and $\mathcal{M}_T$ be the general Markov model on T^r as defined by (2). Then the smallest algebraic variety, i.e. a subset of a real space defined by a finite set of polynomial equations, containing the general Markov model is defined by vanishing of all 3×3 -minors of all the edge flattenings of T^r together with the trivial polynomial equation $\sum_\alpha p_\alpha = 1$.*

Note that the result includes the case of the tripod tree model since in this case each edge flattening of the joint probability table is a 2×4 table so there are no 3×3 minors and hence there are no non-trivial polynomials vanishing on the model.

In an analogous way to the edge flattenings of tables representing probability distributions we can define edge flattenings of $(\kappa_I)_{I \subseteq [n]}$ where $\kappa_\emptyset = 1$ and $\kappa_i = 0$ for all $i \in [n]$ (c.f. Appendix A). Let e be an edge of T inducing a split $(A)(B) \in \Pi_T$ such that $|A| = r$, $|B| = n - r$. Then $\hat{N}_e$ is a $2^r \times 2^{n-r}$ matrix such that for any two subsets $I \subseteq A$, $J \subseteq B$ the element of $\hat{N}_e$ corresponding to the I -th row and the J -th column is κ_{IJ} . Denote by N_e its submatrix given by removing the column and the row corresponding to empty subsets of A and B . Here the labeling for the rows and columns is induced by the ordering of the rows and columns for P_e (c.f. Definition 6.1), i.e. all the subsets of A and B are coded as $\{0,1\}$ -vectors and we introduce the lexicographic order on the vectors with the vector of ones being the last one.

The following result allows us to rephrase the equations in Theorem 6.2 in terms of our new coordinates.

Proposition 6.3. *Let $T = (V, E)$ be a tree and let P be a probability distribution of a vector $X = (X_1, \dots, X_n)$ of binary variables represented by the leaves of T . If $e \in E$ is an edge of T inducing a split $(A_1)(A_2)$ then $\text{rank}(P_e) = 2$ if and only if $\text{rank}(N_e) = 1$.*

Proof. Let $P_e = [p_{\alpha\beta}]$ be the matrix induced by a split $(A_1)(A_2)$. We will show that $\text{rank}(P_e) = \text{rank}(D_e)$ where $D_e = [d_{IJ}]$ is a block diagonal matrix with 1 as the first 1×1 block (i.e. $d_{\emptyset\emptyset} = 1$, $d_{\emptyset J} = 0$, $d_{I\emptyset} = 0$ for all $I \subseteq A_1$, $J \subseteq A_2$) and matrix N_e as the second block. It will then follow that $\text{rank}(P_e) = 2$ if and only if $\text{rank}(N_e) = 1$.

First note that the flattening matrix P_e can be transformed to the flattening of the non-central moments just by adding rows and columns according to (29) and then to the flattening of the central moments $M_e = [\mu_{IJ}]$ such that $I \subseteq A_1$, $J \subseteq A_2$ using (30). It therefore suffices to show that $\text{rank}(M_e) = \text{rank}(D_e)$.

Let $I \subseteq A_1$, $J \subseteq A_2$. Then for each $\pi \in \Pi_{T(IJ)}$ there is at most one block containing elements from both I and J . For otherwise removing e would increase the number of blocks in π by more than one which is not possible. Denote this block by $(I'J')$ where $I' \subseteq I$, $J' \subseteq J$. Note that by construction we have either both I', J' are empty sets if $\pi \geq (A_1)(A_2)$ in $\Pi_{T(IJ)}$ or both $I', J' \neq \emptyset$ otherwise. We can rewrite (32) splitting the blocks

$$(28) \quad \mu_{IJ} = \sum_{\pi \in \Pi_{T(IJ)}} \left(\kappa_{I'J'} \prod_{I \supseteq B \in \pi} \kappa_B \prod_{J \supseteq B \in \pi} \kappa_B \right).$$

We have $d_{I'J'} = \kappa_{I'J'}$ and it can be further rewritten as

$$\mu_{IJ} = \sum_{I' \subseteq I} \sum_{J' \subseteq J} u_{II'} d_{I'J'} v_{J'J}$$

where $u_{II'} = \sum_{\pi \in \Pi_T(I \setminus I')} \prod_{B \in \pi} \kappa_B$ and $v_{J'J} = \sum_{\pi \in \Pi_T(J \setminus J')} \prod_{B \in \pi} \kappa_B$. Setting $u_{II'} = 0$ for $I' \not\subseteq I$, $v_{J'J} = 0$ for $J' \not\subseteq J$ we can write these coefficients in terms of a lower triangular matrix U and an upper triangular matrix V . Since by construction $u_{II} = 1$ for all $I \subseteq A_1$ and $v_{JJ} = 1$ for all $J \subseteq A_2$ we have $\det U = \det V = 1$. Consequently, M_e has the same rank as D_e . $\square$

The proposition shows that the vanishing of all 3×3 minors of all the edge flattenings of P and the trivial invariant $\sum p_\alpha = 1$ are together equivalent to the vanishing all 2×2 minors of all edge flattenings of $\kappa = (\kappa_I)_{I \in [n]_{\geq 2}}$. An immediate corollary follows which gives the equations in (C1) in Theorem (4.6).

Corollary 6.4. *Let $T = (V, E)$ be a trivalent tree. Then the smallest algebraic variety containing $\mathcal{M}_T^\kappa$ is defined by the following set of equations. For each split $(A)(B)$ induced by an edge consider any four (not necessarily disjoint) nonempty sets $I_1, I_2 \subseteq A$, $J_1, J_2 \subseteq B$ and the induced equation $\kappa_{I_1 J_1} \kappa_{I_2 J_2} - \kappa_{I_1 J_2} \kappa_{I_2 J_1} = 0$.*

In [10] Eriksson noted that some of invariants usually prove to be better in discriminating between different tree topologies than the others. His simulations showed that the invariants related to the four-point condition were especially powerful. The binary case we consider in this paper can give some partial understanding of why this might be so. Here, the invariants related to the four-point condition are only those involving second order covariances (c.f. Section 4). Moreover, the estimates of the higher-order moments (or cumulants) are sensitive to outliers and their variance generally grows with the order of the moment. Let $\hat{\mu}$ be a sample estimator of the central moments μ and let f be one of the polynomials in Theorem 6.4 but expressed in terms of the central moments. Then using the delta method we have

$$\text{Var}(f(\hat{\mu})) \simeq \nabla f(\mu)^t \text{Var}(\hat{\mu}) \nabla f(\mu).$$

Consequently, in this loose sense at least, the higher the order of the central moments and hence tree cumulants the higher the variability of we might expect the invariant to exhibit (see [18, Section 4.5]).

7. DISCUSSION

The new coordinate system proposed in [27] provides a better insight into the geometry of phylogenetic tree models with binary observations. The elegant form of the parameterization is useful and has already enabled us to obtain the full geometric description of the model class. One of the interesting implications of this result for phylogenetic tree models is that we could consider different simpler model classes containing the original one in such a way that the whole evolutionary interpretation in terms of the tree topologies remains valid. If we are interested only in the tree we could consider the model defined only by a subsets of constraints in Theorem 4.6 involving only covariances. The price for this reduction is that the conditional independencies induce by the original model do not hold anymore which in turn affects the interpretation of the model. We note that this approach is in a similar spirit to that employed to motivate the MAG model class introduced in [23].

This work has encouraged us to use this reparametrization of this model class to estimate models within Bayesian framework. When the sample proportions lie in the model class then we have already noted that the MLEs are given by formulas in Corollary 11 in [27]. However these sample proportions rarely lie in the model exactly. In a later paper we prove various formal methods for incorporating the semi-algebraic geometry in a model to improve the prior specification of the tree model and hence enhance the estimation of the model parameters.

ACKNOWLEDGEMENTS

Diane Maclagan and John Rhodes contributed substantially to this paper. We would also like to thank Bernd Sturmfels for a stimulating discussion at the early stage of our work and Lior Pachter for pointing out reference [7].

APPENDIX A. CHANGE OF COORDINATES

First we change our coordinates from the raw probabilities $p = [p_\alpha]$ to the non-central moments $\lambda = [\lambda_\alpha]$ for $\alpha = (\alpha_1, \dots, \alpha_n) \in \{0, 1\}^n$, where $\lambda_\alpha = \mathbb{E}(\prod_{i=1}^n X_i^{\alpha_i})$. This is a linear map $f_{p\lambda} : \mathbb{R}^{2^n} \rightarrow \mathbb{R}^{2^n}$ with the determinant equal to one, where the components λ_α of $\lambda = f_{p\lambda}(p)$ are defined by

$$(29) \quad \lambda_\alpha = \sum_{\alpha \leq \beta \leq \mathbf{1}} p_\beta \quad \text{for any } \alpha \in \{0, 1\}^n,$$

where $\mathbf{1}$ denotes here the vector of ones and the sum is over all binary vectors β such that $\alpha \leq \beta \leq \mathbf{1}$ in the sense that $\alpha_i \leq \beta_i \leq 1$ for all $i = 1, \dots, n$. In particular $\lambda_0 = 1$ for all probability distributions. So the image $f_{p\lambda}(\Delta_{2^n-1})$ is contained in the hyperplane defined by $\lambda_0 = 1$.

The linearity of the expectation implies that the central moments can be expressed in terms of non-central moments. Define $\mu_\alpha = \mathbb{E}(\prod_{i=1}^n U_i^{\alpha_i})$, where $U_i = X_i - \mathbb{E}X_i$. Then

$$(30) \quad \mu_\alpha = \sum_{\mathbf{0} \leq \beta \leq \alpha} (-1)^{|\beta|} \lambda_{\alpha-\beta} \prod_{i=1}^n \lambda_{e_i}^{\beta_i} \quad \text{for } \alpha \in \{0, 1\}^n,$$

where $|\beta| = \sum_i \beta_i$. Using these equations we can transform variables from the non-central moments $\lambda = [\lambda_\alpha]$ to another set of variables given by all the means $\lambda_{e_1}, \dots, \lambda_{e_n}$, where $e_1, \dots, e_n$ are standard basis vectors in $\mathbb{R}^n$, and central moments $[\mu_\alpha]$ for $\alpha \in \{0, 1\}$. The polynomial change of coordinates $f_{\lambda\mu} : \mathbb{R}^{2^n} \rightarrow \mathbb{R}^n \times \mathbb{R}^{2^n}$ is an identity on the first n coordinates corresponding to the means $\lambda_{e_1}, \dots, \lambda_{e_n}$ and is defined on the remaining coordinates using the equations (30). Denote $\mathcal{C}_n = (f_{\lambda\mu} \circ f_{p\lambda})(\Delta_{2^n-1})$ which is contained in a subspace of $\mathbb{R}^n \times \mathbb{R}^{2^n}$ given by

$$\mu_0 = 1 \quad \text{and} \quad \mu_{e_1} = \dots = \mu_{e_n} = 0.$$

To simplify notation henceforth we will index moments not with $\{0, 1\}^n$ but with the set of subsets of $[n]$. Here the set $A \subseteq [n]$ is identified with $\alpha \in \{0, 1\}^n$ such that $\alpha_i = 1$ for all $i \in A$ and it is zero elsewhere. In particular for each $i \in [n]$ we write λ_i for λ_{e_i} . The coordinates of $\mathcal{C}_n$ are given by $\lambda_1, \dots, \lambda_n$ together with μ_I for all $I \subseteq [n]$ such that $|I| \geq 2$. Note that the Jacobian of $f_{\lambda\mu} \circ f_{p\lambda} : \Delta_{2^n-1} \rightarrow \mathcal{C}_n$ is constant and equal to one.

The final change of coordinates requires some combinatorics. Let $T = (V, E)$ be a tree with n leaves. A *split induced by* $e \in E$ is a partition of $[n]$ into two non-empty sets induced by removing e from E and restricting $[n]$ to the connected components of the resulting graph. By a *multisplit* we mean any partition $(B_1) \cdots (B_k)$ of the set of leaves induced by removing a subset of the set of edges of T . Each B_i is called a *block* of the partition.

By Π_T we denote the partially ordered set (poset) of all multisplits of the set of leaves induced by edges of T . The poset Π_T has a unique maximal element induced by removing all edges in E and the minimal one with no edges removed which is equal to a single block $[n]$. The maximal element of a lattice is denoted by 1 and the minimal one is denoted by 0.

For any poset Π a *Möbius function* $m_\Pi : \Pi \times \Pi \rightarrow \mathbb{R}$ is defined in such a way that $m_\Pi(x, x) = 1$ for every $x \in \Pi$, $m_\Pi(x, y) = -\sum_{x \leq z < y} m_\Pi(x, z)$ for $x < y$ in Π and it is zero otherwise (c.f. [24, Section 3.7]). Let $W \subset V$, then by $T(W)$ we denote the minimal subtree of T spanned by W . Then $\Pi_{T(W)}$ is the poset of all multisplits of the set of leaves of $T(W)$ induced by edges of $T(W)$. The Möbius function on this poset is denoted by $m_{\Pi_{T(W)}} := m_W$. For Π_T we write $m_{\Pi_T} := m$. We write 0_W and 1_W to denote the minimal and the maximal element of $\Pi_{T(W)}$ respectively.

Consider a map $f_{\mu\kappa} : \mathbb{R}^n \times \mathbb{R}^{2^n} \rightarrow \mathbb{R}^n \times \mathbb{R}^{2^n}$ where the coordinates in the domain are denoted by $\lambda_1, \dots, \lambda_n$ and μ_I for $I \subseteq [n]$ and the coordinates in the image are denoted by $\bar{\mu}_1, \dots, \bar{\mu}_n$ and κ_I for $I \subseteq [n]$. The map is defined by $\bar{\mu}_i = 1 - 2\lambda_i$ for $i = 1, \dots, n$ and

$$(31) \quad \kappa_I = \sum_{\pi \in \Pi_{T(I)}} m_I(0_I, \pi) \prod_{B \in \pi} \mu_B \quad \text{for all } I \subseteq [n],$$

where by convention $\kappa_\emptyset = \mu_\emptyset$. Denote $\mathcal{K}_T = f_{\mu\kappa}(\mathcal{C}_n)$. Note that for any $I \subseteq [n]$ such that $|I| \leq 3$ we have $\kappa_I = \mu_I$. In particular $\mathcal{K}_T$ is contained in the subspace of $\mathbb{R}^n \times \mathbb{R}^{2^n}$ given by

$$\kappa_\emptyset = 1, \quad \kappa_1 = \dots = \kappa_n = 0$$

Therefore the coordinate system on $\mathcal{K}_T$ is given by $\bar{\mu}_1, \dots, \bar{\mu}_n$ and κ_I for $|I| \geq 2$. The map $f_{\mu\kappa} : \mathcal{C}_n \rightarrow \mathcal{K}_T$ is a one-to-one polynomial map with a polynomial inverse $f_{\kappa\mu}$. The exact form of the inverse map is given by the Möbius inversion formula (c.f. [27])

$$(32) \quad \mu_I = \sum_{\pi \in \Pi_{T(I)}} \prod_{B \in \pi} \kappa_B \quad \text{for all } I \subseteq [n], |I| \geq 2,$$

and $\lambda_i = \frac{1}{2}(1 - \bar{\mu}_i)$ for $i = 1, \dots, n$. The Jacobian of $f_{\mu\kappa}$ is constant and equal to 2^n . Note that all $f_{p\lambda}$, $f_{\lambda\mu}$, $f_{\mu\kappa}$, after restriction to Δ_{2^n-1} , $f_{p\lambda}(\Delta_{2^n-1})$ and $\mathcal{C}_n$ respectively, are regular polynomial maps with regular inverses (c.f. [27], Appendix A). This therefore implies that there is a regular one-to-one polynomial map between Δ_{2^n-1} and $\mathcal{K}_T$.

APPENDIX B. THE PROOF OF THE MAIN THEOREM

Let $K \in \mathcal{K}_T$ have coordinates given by $\bar{\mu}_i$ for $i \in [n]$ and κ_I for $I \subseteq [n]$ such that $|I| \geq 2$. Let K^J for $J \subseteq [n]$ denote the projection on coordinates given by $\bar{\mu}_i$ for $i \in J$ and κ_I for $I \subseteq J$ such that $|I| \geq 2$. It follows directly from the definition of $\mathcal{M}_T$ that $K \in \mathcal{M}_T^\kappa$ if and only if $K^I \in \mathcal{M}_{T(I)}^\kappa$ for all $I \subseteq [n]$.

Denote by $\mathcal{M}$ the subset of $\mathcal{K}_T$ defined by constraints in (C1)-(C5). We need to show that $\mathcal{M} = \mathcal{M}_T^\kappa$. Since the rooting is not relevant we choose an arbitrary inner node as the root node. We first show that $\mathcal{M}_T^\kappa \subseteq \mathcal{M}$. Let $K \in \mathcal{M}_T^\kappa$ and hence $K = \psi_T(\omega)$ for some $\omega \in \Omega_T$. The equations in (C1) hold since by construction (c.f. Theorem 6.2) the variety in $\mathcal{K}_T$ defined by the equations contains the image of ψ_T and hence also K . To show that K satisfies (C2) and (C3) consider the projection K^{ijk} for each $i, j, k \in [n]$. By Lemma 1 in [27] $\mathcal{M}_{T(ijk)}^\kappa$ is equal to the tripod tree model. Since $K^{ijk} \in \mathcal{M}_{T(ijk)}^\kappa$ then by Lemma 3.2 (C2) and (C3) must hold. To show that K satisfies (C4) let $i, j \in I$ be such that $\mu_{ij} = 0$. In this case from (7) ω is such that either $\eta_{u,v} = 0$ for some $(u, v) \in E(ij)$ or $\bar{\mu}_{r(ij)}^2 = 1$. In the first case since $E(ij) \subseteq E(I)$ then $\kappa_I = 0$ by (7). In the second case if $r(ij) = r(I)$ then again $\kappa_I = 0$ by (7). If $r(ij) \neq r(I)$ then the edge $(v, r(ij))$ pointing to $r(ij)$ also lies in $E(I)$. By (5) either $\eta_{v,r(ij)} = 0$ in which case we are done or $\bar{\mu}_v^2 = 1$ in which case again there exists an edge in $E(I)$ pointing into v . Since the tree is finite eventually either $\bar{\mu}_{r(I)}^2 = 1$ or $\eta_{u,v} = 0$ for some $(u, v) \in E(I)$. This shows that necessarily $\kappa_I = 0$ and hence K satisfies (C4).

To show that K satisfies (C5) let $i, j, k, l \in [n]$ be the four leaves mentioned in the condition. Let u and v be two inner nodes such that u separates i from j , v separates k from l and $\{u, v\}$ separates $\{i, j\}$ from $\{k, l\}$. In other words u, v are the only inner nodes of degree three in $T(ijkl)$. By Lemma 2 in [27] $T(ijkl)$ gives the same model as the quartet tree with four leaves i, j, k, l and two inner nodes u, v . Moreover, by Remark 2.5, $\mathcal{M}_{T(ijkl)}$ does not depend on the rooting so we can assume that the tree is rooted in u . Since $K^{ijkl} \in \mathcal{M}_{T(ijkl)}$ then for some parameter choices

$$\begin{aligned} \mu_{ik} &= \frac{1}{4}(1 - \bar{\mu}_u^2)\eta_{u,i}\eta_{u,v}\eta_{v,k}, & \mu_{jl} &= \frac{1}{4}(1 - \bar{\mu}_u^2)\eta_{u,j}\eta_{u,v}\eta_{v,l} \\ \mu_{ijk} &= \frac{1}{4}(1 - \bar{\mu}_u^2)\bar{\mu}_u\eta_{u,i}\eta_{u,j}\eta_{u,v}\eta_{v,k}, & \mu_{ikl} &= \frac{1}{4}(1 - \bar{\mu}_u^2)\bar{\mu}_v\eta_{u,i}\eta_{u,v}\eta_{v,k}\eta_{v,l}. \end{aligned}$$

Substitute these equations into (C5). There are then two cases to consider: $\mu_{uv} \geq 0$, $\mu_{uv} < 0$. Laborious but elementary algebra shows that the condition in (C5) is equivalent to (5) applied to $(1 - \bar{\mu}_u^2)\eta_{u,v}$ and hence (C5) holds by definition. Consequently $\mathcal{M}_T^\kappa \subseteq \mathcal{M}$.

We next show that $\mathcal{M} \subseteq \mathcal{M}_T^\kappa$. Let $K \in \mathcal{M}$. We construct a point $\omega_0 \in \mathbb{R}^{|V|+|E|}$ such that $\omega_0 \in \Omega_T$ and $\psi_T(\omega_0) = K$, i.e. ω_0 is such that for all $I \subseteq [n]$ such that $|I| \geq 2$ κ_I can be written in terms of the parameters in ω_0 as in (7).

Case 1: Begin by assuming that K is such that $\mu_{ij} \neq 0$ for all $i, j \in [n]$. We now set squares of values of all the parameters in terms of the observed moments as in Corollary 11 in [27]. We will show that the equations in (7) must hold for their modulus values. Next we will need to ensure there is at least one assignment of signs for a set of parameters such that all (7) hold exactly. Finally we show that the parameter vector ω_0 defined in this way lies in Ω_T .

For each inner node h of T let $i, j, k \in [n]$ be separated by h in T . By (C2) we have that $\mu_{ij}\mu_{ik}\mu_{jk} > 0$ and hence also that $\text{Det}P^{ijk} > 0$. Now set

$$(33) \quad (\bar{\mu}_h^0)^2 = \frac{\mu_{ijk}^2}{\text{Det}P^{ijk}}.$$

We show that (C1), which K satisfies by assumption, implies that the value of $(\bar{\mu}_h^0)^2$ does not depend on the choice of i, j, k . It suffices to show that if k is replaced by another leaf k' such that i, j, k' are separated by h in T then $\frac{\mu_{ijk}^2}{\text{Det}P^{ijk}} = \frac{\mu_{ijk'}^2}{\text{Det}P^{ijk'}}$. Since h has degree

three in T then there exists an edge $e \in E$ inducing a split $(A)(B)$ such that $i, j \in A$ and $k, k' \in B$. From (C1) it follows that

$$(34) \quad \mu_{ik}\mu_{jk'} = \mu_{ik'}\mu_{jk}, \quad \mu_{ijk}\mu_{ik'} = \mu_{ijk'}\mu_{ik}, \quad \mu_{ijk}\mu_{jk'} = \mu_{ijk'}\mu_{jk}$$

and consequently

$$(35) \quad \text{Det} P^{ijk} \mu_{ij} \mu_{ik'} \mu_{jk'} = \text{Det} P^{ijk'} \mu_{ij} \mu_{ik} \mu_{jk}$$

which implies that

$$\frac{\mu_{ijk}^2}{\text{Det} P^{ijk}} = \frac{\mu_{ijk}^2 \mu_{ij} \mu_{ik'} \mu_{jk'}}{\text{Det} P^{ijk} \mu_{ij} \mu_{ik'} \mu_{jk'}} = \frac{\mu_{ijk'}^2 \mu_{ij} \mu_{ik} \mu_{jk}}{\text{Det} P^{ijk'} \mu_{ij} \mu_{ik} \mu_{jk}} = \frac{\mu_{ijk'}^2}{\text{Det} P^{ijk'}}$$

as required.

For terminal edges (v, i) of T such that $i \in [n]$ let $j, k \in [n]$ be any two leaves of T such that v separates i, j, k . Set

$$(36) \quad (\eta_{v,i}^0)^2 = \frac{\text{Det} P^{ijk}}{\mu_{jk}^2}.$$

As in the previous case it is straightforward to check that given (C1) this value does not depend on the choice of j, k . Without loss assume that instead of k we have k' and v separates i, j, k' in T . Since there exists an edge split such that i, j and k, k' are in different blocks we have (34) and (34) and consequently

$$\frac{\text{Det} P^{ijk}}{\mu_{jk}^2} = \frac{\mu_{ik} \text{Det} P^{ijk}}{\mu_{ik'} \mu_{jk'} \mu_{jk}} = \frac{\text{Det} P^{ijk'}}{\mu_{jk'}^2}.$$

For inner edges $(u, v) \in E$ let $i, j, k, l \in [n]$ be any four leaves such that u separates i from j , v separates k from l and $\{u, v\}$ separates $\{i, j\}$ from $\{k, l\}$. Set

$$(37) \quad (\eta_{u,v}^0)^2 = \frac{\mu_{il}^2}{\mu_{ij}^2} \frac{\text{Det} P^{ijk}}{\text{Det} P^{ikl}}$$

which is well-defined since μ_{ij}^2 and $\text{Det} P^{ikl}$ are strictly positive. We now show that this value does not depend on the choice of i, j, k, l . By symmetry it suffices to show that we obtain the same value if instead of l we took another leaf l' such that u, v are the only degree three nodes in $T(ijkl')$. Since v has degree three then there must exist an inner edge separating i, j, k from l, l' . From (C1) it follows that

$$\mu_{il'} \mu_{kl'} \text{Det} P^{ikl} = \mu_{il} \mu_{kl} \text{Det} P^{ikl'}, \quad \mu_{il} \mu_{kl'} = \mu_{il'} \mu_{kl}$$

and hence

$$\frac{\mu_{il}^2}{\mu_{ij}^2} \frac{\text{Det} P^{ijk}}{\text{Det} P^{ikl}} = \frac{\mu_{il'} \mu_{kl'} \mu_{il}^2}{\mu_{il'} \mu_{kl'} \mu_{ij}^2} \frac{\text{Det} P^{ijk}}{\text{Det} P^{ikl}} = \frac{\mu_{il'}^2}{\mu_{ij}^2} \frac{\text{Det} P^{ijk}}{\text{Det} P^{ikl'}}$$

as required.

We now show that the modulus of equations (7) hold. First consider the case $I = \{i, j\}$. Label the inner nodes of $E(ij)$ by $v_1, \dots, v_k$ beginning from the node adjacent to i . For each $s = 1, \dots, k$ let i_s denote a leaf such that v_s separates i, j, i_s in T . We assume that the root $r(ij)$ of this path is in v_1 . The analysis is the same for any other rooting by Remark 2.5. We now proceed to check that

$$(38) \quad \begin{aligned} \mu_{ij}^2 &= \left(\frac{1}{4} (1 - (\bar{\mu}_{r(ij)}^0)^2) \right)^2 \prod_{(u,v) \in E(ij)} (\eta_{u,v}^0)^2 = \\ &= \left(\frac{1}{4} (1 - (\bar{\mu}_{v_1}^0)^2) \right)^2 (\eta_{v_1,u}^0)^2 \left(\prod_{s=2}^k (\eta_{v_{s-1}, v_s}^0)^2 \right) (\eta_{v_k, v}^0)^2. \end{aligned}$$

Since v_1 separates i, j, i_1 by construction, from (33) we therefore have

$$\frac{1}{4} (1 - (\bar{\mu}_{v_1}^0)^2) = \frac{\mu_{ij} \mu_{i_1} \mu_{j i_1}}{\text{Det}(P^{ij i_1})}.$$

Now substitute this equation and all the set values in (36), (37) into the right hand side of (38). Use the fact that v_k separates i, j, i_k in T and i_{s-1}, i_s are the only degree three nodes in $T(ii_{s-1}ji_s)$. Since (v_1, i) and (v_k, j) are the only terminal edges we obtain

$$(39) \quad \left(\frac{\mu_{ij}\mu_{ii_1}\mu_{ji_1}}{\text{Det}(P^{iji_1})} \right)^2 \cdot \frac{\text{Det}P^{iji_1}}{\mu_{ji_1}^2} \cdot \left(\prod_{s=2}^k \frac{\mu_{ii_s}^2}{\mu_{ii_{s-1}}^2} \frac{\text{Det}P^{iji_{s-1}}}{\text{Det}P^{iji_s}} \right) \cdot \frac{\text{Det}P^{iji_k}}{\mu_{ji_k}^2}$$

It can now be checked that all the expressions with hyperdeterminants cancel out and the formula reduces to μ_{ij}^2 as required.

Now we need to show that for every $I = \{i, j, k\}$

$$(40) \quad \mu_{ijk}^2 = \left(\frac{1}{4}(1 - \bar{\mu}_{r(ijk)}^0)^2 \right)^2 (\bar{\mu}_w^0)^2 \prod_{(u,v) \in E(ijk)} (\eta_{u,v}^0)^2,$$

where by w we denote the node separating i, j and k . Assume that $T(ijk)$ is rooted somewhere on the path between i and j . Using (38) the right hand side of (40) can be rewritten as

$$(41) \quad \mu_{ij}^2 (\bar{\mu}_w^0)^2 \prod_{(u,v) \in E(wk)} (\eta_{u,v}^0)^2.$$

Number the degree three nodes in $E(wk)$ by $v_1, \dots, v_l$ and let i_s denote a leaf such that the inner nodes of $T(ijk i_s)$ of degree three are exactly v_{s-1} and v_s , where $v_0 = w$. By an exactly analogous argument as in the case above we obtain

$$(42) \quad \prod_{(u,v) \in E(wk)} (\eta_{u,v}^0)^2 = \frac{\mu_{ii_1}^2}{\mu_{ij}^2} \frac{\text{Det}P^{ijk}}{\text{Det}P^{iki_1}} \cdot \left(\prod_{s=2}^l \frac{\mu_{i_{s-1}i_s}^2}{\mu_{i_{s-2}i_{s-1}}^2} \frac{\text{Det}P^{i_{s-2}i_{s-1}i_s}}{\text{Det}P^{i_{s-1}i_s i_k}} \right) \frac{\text{Det}P^{i_{l-1}i_l i_k}}{\mu_{i_{l-1}i_l}^2},$$

where $i_0 = i$. It can be easily checked that all the hyperdeterminants apart from the term $\text{Det}P^{ijk}$ cancel out. Moreover all the covariances apart from μ_{ij}^{-2} cancel out as well and hence (42) is equal to $\frac{\text{Det}P^{ijk}}{\mu_{ij}^2}$. Now using the definition of $(\bar{\mu}_w^0)^2$ in (33) it can be easily checked that (41) is equal to μ_{ijk}^2 as required.

So far we have confirmed only that the squares of parameters in ω_0 satisfy required equations at least for the tree cumulants up to the third order. Next, we show that there exists a consistent choice of signs for these parameters such that the equations are satisfied exactly. Let $\sigma(i, j) = \text{sgn}(\mu_{ij})$. Since by assumption $\mu_{ij} \neq 0$ for all $i, j \in [n]$ then the conditions in (C2) imply that $\sigma(i, j)\sigma(i, k)\sigma(j, k) = 1$ for all triples $i, j, k \in [n]$. Hence by Lemma 4.4 there exists a choice $s(u, v) \in \{-1, +1\}$ for all $(u, v) \in E$ such that $\sigma(i, j) = \prod_{(u,v) \in E(ij)} s(u, v)$ for all $i, j \in [n]$. For any two nodes $k, l \in V$ we define $s(k, l) = \prod_{(u,v) \in E(kl)} s(u, v)$. A choice of signs for the parameters can be obtained as follows. For each edge $(u, v) \in E$ we set $\text{sgn}(\eta_{u,v}^0) = s(u, v)$ and for each inner node v we set $\text{sgn}(\bar{\mu}_v^0) = \text{sgn}(\mu_{ijk})s(v, i)s(v, j)s(v, k)$ where i, j, k are any three leaves of T separated by v .

Assume now that the choice of the signs of the parameters, induced by $s(u, v)$ for $(u, v) \in E$, has been made. This choice of signs gives

$$(43) \quad \bar{\mu}_v^0 = s(v, i)s(v, j)s(v, k) \frac{\mu_{ijk}}{\sqrt{\text{Det}P^{ijk}}},$$

$$(44) \quad \eta_{v,i}^0 = s(v, i) \frac{\sqrt{\text{Det}P^{ijk}}}{|\mu_{jk}|},$$

$$(45) \quad \eta_{u,v}^0 = s(u, v) \left| \frac{\mu_{il}}{\mu_{ij}} \right| \sqrt{\frac{\text{Det}P^{ijk}}{\text{Det}P^{ikl}}}.$$

Note that in particular with this choice of signs $\text{sgn}(\eta_{u,v}^0) = s(u, v)$ for all $(u, v) \in E$ and $\text{sgn}(\bar{\mu}_v^0) = \text{sgn}(\mu_{ijk}) \prod_{(u,v) \in E(ijk)} s(u, v)$. For μ_{ij} and μ_{ijk} we have shown that modulus

of (7) hold. It suffices to check that the signs match. But this follows directly from the construction. Indeed, proving (38) and we have shown that

$$|\mu_{ij}| = \frac{1}{4}(1 - (\bar{\mu}_{r(ij)}^0)^2) \prod_{(u,v) \in E(ij)} |\eta_{u,v}^0|.$$

Now multiply both sides by $s(i, j) = \prod_{(u,v) \in E(ij)} s(u, v)$ to get

$$(46) \quad \mu_{ij} = s(i, j)|\mu_{ij}| = \frac{1}{4}(1 - (\bar{\mu}_{r(ij)}^0)^2) \prod_{(u,v) \in E(ij)} s(u, v)|\eta_{u,v}^0| =$$

$$(47) \quad = \frac{1}{4}(1 - (\bar{\mu}_{r(ij)}^0)^2) \prod_{(u,v) \in E(ij)} \eta_{u,v}^0.$$

Similarly from (40) we have that

$$|\mu_{ijk}| = \frac{1}{4}(1 - (\bar{\mu}_{r(ijk)}^0)^2)|\bar{\mu}_w^0| \prod_{(u,v) \in E(ijk)} |\eta_{u,v}^0|.$$

Multiply both sides by $\text{sgn}(\mu_{ijk})$ and use the fact that $(\prod_{(u,v) \in E(ijk)} s(u, v))^2 = 1$ to get

$$\begin{aligned} \mu_{ijk} &= \frac{1}{4}(1 - (\bar{\mu}_{r(ijk)}^0)^2) \left(|\bar{\mu}_w^0| \text{sgn}(\mu_{ijk}) \prod_{(u,v) \in E(ijk)} s(u, v) \right) \prod_{(u,v) \in E(ijk)} s(u, v)|\eta_{u,v}^0| = \\ &= \frac{1}{4}(1 - (\bar{\mu}_{r(ijk)}^0)^2) \bar{\mu}_w^0 \prod_{(u,v) \in E(ijk)} \eta_{u,v}^0 \end{aligned}$$

as desired.

We now show (7) for $|I| \geq 4$ by induction. Let $(u, v) \in E$ be any edge splitting I into two subsets I_1 and I_2 such that $|I_1|, |I_2| \geq 2$ and u is the node closer to I_1 . Let $i \in I_1$ and $j \in I_2$ then by (C1)

$$\kappa_{I_1 I_2} = \frac{\kappa_{I_1 j} \kappa_{i I_2}}{\kappa_{ij}}.$$

By induction we can assume that $\kappa_{I_1 j}$, $\kappa_{i I_2}$ and κ_{ij} have form as in (7). Moreover,

$$\begin{aligned} \frac{\prod_{(u,v) \in E(iI_2)} \eta_{u,v} \prod_{(u,v) \in E(I_1 j)} \eta_{u,v}}{\prod_{(u,v) \in E(ij)} \eta_{u,v}} &= \prod_{(u,v) \in E(I)} \eta_{u,v}, \\ \prod_{h \in N(iI_2)} \bar{\mu}_h^{\deg h - 2} &= \prod_{h \in N(vI_2)} \bar{\mu}_h^{\deg h - 2}, \\ \prod_{h \in N(I_1 j)} \bar{\mu}_h^{\deg h - 2} &= \prod_{h \in N(I_1 u)} \bar{\mu}_h^{\deg h - 2}. \end{aligned}$$

Using this we can write

$$(48) \quad \kappa_{I_1 I_2} = \frac{1}{4} \frac{(1 - \bar{\mu}_{r(iI_2)}^2)(1 - \bar{\mu}_{r(I_1 j)}^2)}{(1 - \bar{\mu}_{r(ij)}^2)} \prod_{h \in N(I)} \bar{\mu}_h^{\deg h - 2} \prod_{(u,v) \in E(I)} \eta_{u,v}.$$

The root of $T(I)$ is either in $T(I_1 u)$ or in $T(vI_2)$. In the first case $r(I_1 j) = r(I)$ and $r(iI_2) = r(ij)$. In the second case $r(I_1 j) = r(ij)$ and $r(iI_2) = r(I)$. Hence in both cases

$$\frac{(1 - \bar{\mu}_{r(iI_2)}^2)(1 - \bar{\mu}_{r(I_1 j)}^2)}{(1 - \bar{\mu}_{r(ij)}^2)} = (1 - \bar{\mu}_{r(I)}^2)$$

and (48) has the required form given by (32). It follows that $K = \psi_T(\omega_0)$.

It now remains to show that the parameters defined in (43), (44) and (45) define a parameter vector ω_0 which lies in Ω_T . Since by (C2) $\mu_{ijk}^2 \leq \text{Det} P^{ijk}$ for all $i, j, k \in [n]$ it follows that for all inner nodes h we have $\bar{\mu}_h^0 \in [-1, 1]$ as required. For a terminal edge (v, i) consider the marginal model induced by $T(ijk)$, where j, k are any two leaves such

that v separates i, j, k in T . From Lemma 3.2 constraints (C2) and (C3) imply that $\eta_{v,i}$ is a valid parameter. To show that (45) satisfies (5) we write

$$(1 - (\bar{\mu}_u^0)^2) \eta_{u,v}^0 = \frac{4\mu_{ij}\mu_{ik}\mu_{jk}}{\sqrt{\text{Det}P^{ijk}\text{Det}P^{ikl}}} s(u, v) \left| \frac{\mu_{il}}{\mu_{ij}} \right|.$$

We now substitute this together with the expressions for $\bar{\mu}_u^0$ and $\bar{\mu}_v^0$ given by (43) into (5). First assume $s(u, v) = 1$ then $s(u, k) = s(v, k)$, $s(v, i) = s(u, i)$ and (5) becomes

$$\frac{4\mu_{ij}\mu_{ik}\mu_{jk}}{\sqrt{\text{Det}P^{ijk}\text{Det}P^{ikl}}} \left| \frac{\mu_{il}}{\mu_{ij}} \right| \leq \left(1 \pm s(u, j) \frac{\mu_{ijk}}{\sqrt{\text{Det}P^{ijk}}} \right) \left(1 \mp s(v, l) \frac{\mu_{ikl}}{\sqrt{\text{Det}P^{ikl}}} \right).$$

The left hand side is equal to $\frac{4\mu_{ik}\mu_{jk}\mu_{il}}{\sqrt{\text{Det}P^{ijk}\text{Det}P^{ikl}}} s(j, l) = \frac{4\mu_{ik}^2\mu_{jl}}{\sqrt{\text{Det}P^{ijk}\text{Det}P^{ikl}}} s(j, l)$. Now multiply both sides by $|\mu_{jl}| \sqrt{\text{Det}P^{ijk}\text{Det}P^{ikl}}$ to get

$$4\mu_{ik}^2\mu_{jl}^2 \leq \left(\sqrt{\mu_{jl}^2 \text{Det}P^{ijk}} \pm s(u, l) \mu_{jl} \mu_{ijk} \right) \left(\sqrt{\text{Det}P^{ikl}} \mp s(v, l) \mu_{ikl} \right).$$

However, $s(u, l) = s(v, l)$ hence this is satisfied by (C5). It is easily calculated that the case $s(u, v) = -1$ leads to the same constraint. This finishes the proof in Case 1 when K is such that $\mu_{ij} \neq 0$ for all $i, j \in [n]$.

Case 2: For the general case let $K \in \mathcal{M}$ be a tree cumulant and let $\Sigma = [\mu_{ij}] \in \mathbb{R}^{n \times n}$ be the matrix of all covariances between the leaves. We say that an edge $e \in E$ is *isolated relative to K* if $\mu_{ij} = 0$ for all $i, j \in [n]$ such that $e \in E(ij)$. By $\widehat{E} \subseteq E$ we denote the set of all edges of T which are isolated relative to K . By $\widehat{T} = (V, E \setminus \widehat{E})$ we denote the forest obtained from T by removing edges in $\widehat{E}$ and we call it the K -forest. We define relations on $\widehat{E}$ and $E \setminus \widehat{E}$. For two edges e, e' with either $\{e, e'\} \subset \widehat{E}$ or $\{e, e'\} \subset E \setminus \widehat{E}$ write $e \sim e'$ if either $e = e'$ or e and e' are adjacent and all the edges that are incident with both e and e' are isolated relative to K . Let us now take the transitive closure of $\sim$ restricted to pairs of edges in $\widehat{E}$ to form an equivalence relation on $\widehat{E}$. This transitive closure is constructed as follows. Consider a graph with nodes representing elements of $\widehat{E}$ and put an edge between e, e' whenever $e \sim e'$. Then the equivalence classes correspond to connected components of this graph. Similarly, take the transitive closure of $\sim$ restricted to the pairs of edges in $E \setminus \widehat{E}$ to form an equivalence relation in $E \setminus \widehat{E}$. We will let $[\widehat{E}]$ and $[E \setminus \widehat{E}]$ denote the set of equivalence classes of $\widehat{E}$ and $E \setminus \widehat{E}$ respectively (for details see Section 5 in [27]).

Again we show that there exists $\omega_0 \in \Omega_T$ such that $\psi_T(\omega_0) = K$. Set $\eta_{u,v}^0 = 0$ for all $(u, v) \in \widehat{E}$ and $\bar{\mu}_v^0 = 0$ for all inner nodes of T with degree zero in $\widehat{T}$. It then follows that $(1 - \bar{\mu}_u^2) \eta_{u,v} = 0$ satisfies (5) for all $(u, v) \in \widehat{E}$ and $\bar{\mu}_v^0 \in [-1, 1]$ for all $v \in \widehat{V}$ and hence these parameters satisfy constraints defining Ω_T . If $I \subseteq [n]$ is such that $E(I) \cap \widehat{E} \neq \emptyset$ then $\kappa_I = 0$ by (C4). Hence in this case we can assert that

$$\kappa_I = \frac{1}{4} (1 - (\bar{\mu}_{r(I)}^0)^2) \prod_{v \in N(I)} (\bar{\mu}_v^0)^{\deg(v)-2} \prod_{(u,v) \in E(I)} \eta_{u,v}^0$$

simply because both sides of this equation are zero. By Lemma 9 (iv) in [27] every connected component of $\widehat{T}$ is a subtree which is either an inner node or a tree with the set of leaves contained in $[n]$. Denote the connected subtrees which are not inner nodes by $T_1, \dots, T_k$ and their sets of leaves by $[n_l]$ for $l = 1, \dots, k$. For every $l = 1, \dots, k$ and all $i, j \in [n_l]$ we have that $\mu_{ij} \neq 0$. Hence for each T_l applying Case 1 we have $K^{[n_l]} \in \mathcal{M}_{T_l}$. If $I \subseteq [n]$ is such that $E(I) \cap \widehat{E} = \emptyset$ then $I \subseteq [n_l]$ for some $l = 1, \dots, k$. Since $K^{[n_l]} \in \mathcal{M}_{T_l}$ then there exists a choice of parameters such that κ_I can be written as (7). Consequently $K \in \mathcal{M}_T$ and we are finished. $\square$

REFERENCES

- [1] E. S. ALLMAN AND J. A. RHODES, *Phylogenetic ideals and varieties for the general Markov model*, Adv. in Appl. Math., 40 (2008), pp. 127–148.
- [2] V. AUVRAY, P. GEURTS, AND L. WEHENKEL, *A Semi-Algebraic Description of Discrete Naive Bayes Models with Two Hidden Classes*, in Proc. Ninth International Symposium on Artificial Intelligence and Mathematics, Fort Lauderdale, Florida, Jan 2006.
- [3] S. BASU, R. POLLACK, AND M. ROY, *Algorithms in Real Algebraic Geometry*, Springer, 2003.
- [4] P. BUNEMAN, *A note on the metric properties of trees*, J. Combinatorial Theory Ser. B, 17 (1974), pp. 48–50.
- [5] M. CASANELLAS AND J. FERNANDEZ-SANCHEZ, *Performance of a New Invariants Method on Homogeneous and Nonhomogeneous Quartet Trees*, Molecular Biology and Evolution, 24 (2007), p. 288.
- [6] J. CAVENDER AND J. FELSENSTEIN, *Invariants of phylogenies in a simple case with discrete states*, Journal of Classification, 4 (1987), pp. 57–71.
- [7] J. A. CAVENDER, *Letter to the editor*, Molecular Phylogenetics and Evolution, 8 (1997), pp. 443 – 444.
- [8] M. DRTON, B. STURMFELS, AND S. SULLIVANT, *Algebraic factor analysis: tetrads, pentads and beyond*, Probab. Theory Related Fields, 138 (2007), pp. 463–493.
- [9] M. DRTON AND S. SULLIVANT, *Algebraic Statistical Models*, Statistica Sinica, 17 (2007), pp. 1273–1297.
- [10] N. ERIKSSON, *Using invariants for phylogenetic tree construction*, vol. 149 of The IMA Volumes in Mathematics and its Applications, Springer, 2007, pp. 89–108.
- [11] N. ERIKSSON, K. RANESTAD, B. STURMFELS, AND S. SULLIVANT, *Phylogenetic algebraic geometry*, in Projective varieties with unexpected properties, Walter de Gruyter GmbH & Co. KG, Berlin, 2005, pp. 237–255.
- [12] L. GARCIA, M. STILLMAN, AND B. STURMFELS, *Algebraic geometry of Bayesian networks*, J. Symbolic Comput, 39 (2005), pp. 331–355.
- [13] I. GELFAND, M. KAPRANOV, AND A. ZELEVINSKY, *Discriminants, Resultants, and Multidimensional Determinants*, Birkhäuser, 1994.
- [14] Z. GILULA, *Singular value decomposition of probability matrices: Probabilistic aspects of latent dichotomous variables*, Biometrika, 66 (1979), pp. 339–344.
- [15] J. LAKE, *A rate-independent technique for analysis of nucleic acid sequences: evolutionary parsimony*, 1987.
- [16] P. LAZARSFELD AND N. HENRY, *Latent structure analysis*, Houghton, Mifflin, New York, 1968.
- [17] F. MATSEN, *Fourier transform inequalities for phylogenetic trees*, Computational Biology and Bioinformatics, IEEE/ACM Transactions on, 6 (2009), pp. 89–95.
- [18] P. MCCULLAGH, *Tensor methods in statistics*, Monographs on Statistics and Applied Probability, Chapman & Hall, London, 1987.
- [19] J. PEARL AND M. TARSI, *Structuring causal trees*, J. Complexity, 2 (1986), pp. 60–77. Complexity of approximately solved problems (Morningside Heights, N.Y., 1985).
- [20] G.-C. ROTA AND J. SHEN, *On the combinatorics of cumulants*, J. Combin. Theory Ser. A, 91 (2000), pp. 283–304. In memory of Gian-Carlo Rota.
- [21] C. SEMPLE AND M. STEEL, *Phylogenetics*, vol. 24 of Oxford Lecture Series in Mathematics and its Applications, Oxford University Press, Oxford, 2003.
- [22] R. SETTIMI AND J. Q. SMITH, *On the Geometry of Bayesian Graphical Models with Hidden Variables*, in UAI, G. F. Cooper and S. Moral, eds., Morgan Kaufmann, 1998, pp. 472–479.
- [23] P. SPIRITES, T. RICHARDSON, AND C. MEEK, *Heuristic greedy search algorithms for latent variable models*, in Proceedings of AI & STAT’97, Citeseer, pp. 481–488.
- [24] R. P. STANLEY, *Enumerative combinatorics. Volume I*, no. 49 in Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2002.
- [25] M. STEEL AND B. FALLER, *Markovian log-supermodularity, and its applications in phylogenetics*, Applied Mathematics Letters, (2009).
- [26] B. STURMFELS AND S. SULLIVANT, *Toric Ideals of Phylogenetic Invariants*, Journal of Computational Biology, 12 (2005), pp. 204–228.
- [27] P. ZWIERNIK AND J. Q. SMITH, *Tree-cumulants and the identifiability of Bayesian tree models*, [arXiv:1004.4360](https://arxiv.org/abs/1004.4360), 2010. submitted.

PIOTR ZWIERNIK, UNIVERSITY OF WARWICK, DEPARTMENT OF STATISTICS, CV7AL, COVENTRY,
UK.

E-mail address: `p.w.zwiernik@warwick.ac.uk`

JIM Q. SMITH, UNIVERSITY OF WARWICK, DEPARTMENT OF STATISTICS, CV7AL, COVENTRY,
UK.